

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 498/04, A61K 31/44, C07D 513/04, 517/04, 513/14 // (C07D 498/04, 263:00, 221:00) (C07D 513/04, 277:00, 221:00) (C07D 517/04, 293:00, 221:00) (C07D 513/14, 277:00, 221:00, 221:00)	A1	(11) International Publication Number: WO 00/06577 (43) International Publication Date: 10 February 2000 (10.02.00)
(21) International Application Number: PCT/US99/17027 (22) International Filing Date: 28 July 1999 (28.07.99) (30) Priority Data: 60/094,346 28 July 1998 (28.07.98) US 09/361,544 27 July 1999 (27.07.99) US (71) Applicant: 3M INNOVATIVE PROPERTIES COMPANY [US/US]; 3M Center, P.O. Box 33427, Saint Paul, MN 55133-3427 (US). (72) Inventors: GERSTER, John, F.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). LINDSTROM, Kyle, J.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). MARSZALEK, Gregory, J.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). MERRILL, Bryon, A.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). MICKELSON, John, W.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US). RICE, Michael, J.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US).		(74) Agents: HOWARD, MarySusan et al.; 3M Innovative Properties Company, Office of Intellectual Property Counsel, P.O. Box 33427, Saint Paul, MN 55133-3427 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: OXAZOLO, THIAZOLO AND SELENAZOLO [4,5-c]-QUINOLIN-4-AMINES AND ANALOGS THEREOF (57) Abstract Thiazolo-, oxazolo- and selenazolo[4,5-c]quinoline-4-amines and analogs thereof are described including methods of manufacture and the use of novel intermediates. The compounds are immunomodulators and induce cytokine biosynthesis, including interferon and/or tumor biosynthesis, necrosis factor, and inhibit the T-helper-type 2 immune response. The compounds are further useful in the treatment of viral and neoplastic diseases.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

OXAZOLO, THIAZOLO AND SELENAZOLO
[4,5-c]-QUINOLIN-4-AMINES AND ANALOGS THEREOF

Field of the Invention

5 This invention relates to oxazolo, thiazolo and selenazolo [4,5-c]-quinolin-,
tetrahydroquinolin-4-amines and hetero analogs thereof, and to intermediates used in their
preparation. The invention also relates to pharmaceutical compositions containing the
above compounds as well as the use of these compounds as immunomodulators and for
inducing cytokine biosynthesis, including interferon- α biosynthesis and/or tumor necrosis
10 factor- α biosynthesis.

Background of the Invention and Related Prior Art


 The first reliable report on the 1H-imidazo[4,5-c]quinoline ring system, Backman
et al., J. Org. Chem. 15, 1278-1284 (1950) describes the synthesis of 1-(6-methoxy-8-
15 quinoliny)-2-methyl-1H-imidazo[4,5-c]quinoline for possible use as an antimalarial agent.
Subsequently, syntheses of various substituted 1H-imidazo[4,5-c] quinolines have been
reported. For example, Jain et al., J. Med. Chem. 11, pp. 87-92 (1968), has synthesized
the compound 1-[2-(4-piperidyl)ethyl]-1H-imidazo[4,5-c]quinoline as a possible
anticonvulsant and cardiovascular agent. Also, Baranov et al., Chem. Abs. 85, 94362
20 (1976), and Berenyi et al., J. Heterocyclic Chem. 18, 1537-1540 (1981), have reported
certain 2-oxoimidazo[4,5-c]quinolines.

 Following the above report, 1H-imidazo[4,5-c]quinolin-4-amines and 1- and 2-
substituted derivatives thereof were found to be useful as antiviral agents, bronchodilators
and immunomodulators. These are described in U.S. Patent Nos. 4,689,338; 4,698,348;
25 4,929,624; 5,037,986; 5,266,675; 5,268,376; 5,346,905; 5,389,640; 5,605,899; 5,352,784;
5,446,153; and 5,482,936. Shen et al., U.S. Patent Nos. 4,038,396 and 4,131,677, describe
certain oxazolo-and thiazolopyridines as having antiinflammatory, analgesic, and
antipyretic properties.

30

Summary of the Invention

The present invention provides compounds of the Formula I



(I)

wherein:

5 **R₁** is selected from the group consisting of oxygen, sulfur and selenium;

R₂ is selected from the group consisting of

- hydrogen;
- alkyl;
- alkyl-OH;
- 10 -haloalkyl;
- alkenyl;
- alkyl-X-alkyl;
- alkyl-X-alkenyl;
- alkenyl-X-alkyl;
- 15 -alkenyl-X-alkenyl;
- alkyl-N(R₅)₂;
- alkyl-N₃;
- alkyl-O-C(O)-N(R₅)₂;
- heterocyclyl;
- 20 -alkyl-X-heterocyclyl;
- alkenyl-X-heterocyclyl;
- aryl;
- alkyl-X-aryl;
- alkenyl-X-aryl;
- 25 -heteroaryl;
- alkyl-X-heteroaryl; and

-alkenyl-X-heteroaryl;

R_3 and R_4 are each independently:

-hydrogen;

-X-alkyl;

5 -halo;

-haloalkyl;

-N(R_5)₂;

or when taken together, R_3 and R_4 form a fused


aromatic, heteroaromatic, cycloalkyl or heterocyclic ring;

10 X is selected from the group consisting of -O-, -S-, -NR₅-, -C(O)-, -C(O)O-,
-OC(O)-, and a bond; and

each R_5 is independently H or C₁₋₈alkyl;

with the proviso that when R_1 is sulfur, R_3 is not -NH₂; or a pharmaceutically acceptable salt thereof.

15 As a second aspect, the present invention provides pharmaceutical compositions containing a therapeutically effective amount of a compound of Formula I(a) and a pharmaceutically acceptable vehicle:



I(a)

20

wherein:

R_1 is selected from the group consisting of oxygen, sulfur and selenium;

R_2 is selected from the group consisting of

-hydrogen;

25 -alkyl;

- alkyl-OH;
- haloalkyl;
- alkenyl;
- alkyl-X-alkyl;
- 5 -alkyl-X-alkenyl;
- alkenyl-X-alkyl;
- alkenyl-X-alkenyl;
- alkyl-N(R₅)₂;
- alkyl-N₃;
- 10 -alkyl-O-C(O)-N(R₅)₂;
- heterocyclyl;
- alkyl-X-heterocyclyl;
- alkenyl-X-heterocyclyl;
- aryl;
- 15 -alkyl-X-aryl;
- alkenyl-X-aryl;
- heteroaryl;
- alkyl-X-heteroaryl; and
- alkenyl-X-heteroaryl;

20 **R₃ and R₄ are each independently:**

- hydrogen;
- X-alkyl;
- halo;
- haloalkyl;
- 25 -N(R₅)₂;


or when taken together, **R₃ and R₄** form a fused aromatic, heteroaromatic, cycloalkyl or heterocyclic ring;

X is selected from the group consisting of -O-, -S-, -NR₅-, -C(O)-, -C(O)O-, -OC(O)-, and a bond; and

30 each **R₅** is independently H or C₁₋₈alkyl; or a pharmaceutically acceptable salt thereof.

The compounds of Formula I(a) are useful in inducing the biosynthesis of certain cytokines in animals, including humans. Cytokines that may be induced by the compounds of the invention include but are not limited to, interferons, particularly
5 interferon- α , and tumor necrosis factor- α . The invention therefore also provides a method of inducing cytokine biosynthesis in an animal by administering to the animal an effective amount of a composition comprising a compound of Formula I(a). Because of their ability to induce cytokine biosynthesis the compounds of the invention are useful in the treatment of a variety of conditions, including viral and neoplastic diseases, and the invention further
10 provides a method of treating such conditions in a subject by administering a therapeutically effective amount of a composition comprising a compound of Formula I(a) to the subject.

As yet another aspect, the present invention provides intermediate compounds of Formula II



wherein:

R_1 is selected from the group consisting of oxygen, sulfur and selenium;

20 R_2 is selected from the group consisting of

-hydrogen;

-alkyl;

-alkyl-OH;

-haloalkyl;

25 -alkenyl;

-alkyl-X-alkyl;

- 5 -alkyl-X-alkenyl;
 -alkenyl-X-alkyl;
 -alkenyl-X-alkenyl;
 -alkyl-N(R₅)₂;
 -alkyl-N₃;
 -alkyl-O-C(O)-N(R₅)₂;
 -heterocyclyl;
 -alkyl-X-heterocyclyl;
 -alkenyl-X-heterocyclyl;
10 -aryl;
 -alkyl-X-aryl;
 -alkenyl-X-aryl;
 -heteroaryl;
 -alkyl-X-heteroaryl;
15 -alkenyl-X-heteroaryl;
 -SO₂CH₃; and
 -CH₂-O-C(O)-CH₃;

R₃ and R₄ are each independently:

- 20 -hydrogen;
 -X-alkyl;
 -halo;
 -haloalkyl;
 -N(R₅)₂;
 or when taken together, **R₃ and R₄** form a fused
25 aromatic, heteroaromatic, cycloalkyl or heterocyclic ring;

X is selected from the group consisting of -O-, -S-, -NR₅-, -C(O)-, -C(O)O-,
and a bond; and

each **R₅** is independently H or C₁₋₈alkyl.


Detailed Description of the Invention

This invention includes compounds of Formula I, pharmaceutical compositions containing compounds of Formula I(a) and therapeutic methods using compounds of Formula I(a) as well as intermediate compounds of Formula II that are used to prepare the compounds of Formulae I and I(a).

The terms "alkyl" and "alkenyl" as used herein refer to a straight or branched hydrocarbon group, or a cyclic group (i.e., cycloalkyl and cycloalkenyl) that contains from 1 to 20, preferably 1 to 10, more preferably 1 to 8 carbon atoms, unless otherwise specified. Typical alkyl groups are, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. Exemplary cyclic groups include cyclopropyl, cyclopentyl, cyclohexyl, cyclohexenyl and adamantyl. The prefix "alk," when used, e.g. for "alkoxy" and the like, also has the same meaning.

The term "aryl" refers to a carbocyclic aromatic ring or ring system. The aryl group is preferably a six-membered ring, such as phenyl, or an aromatic polycyclic ring system, such as naphthyl. The most preferred aryl group is phenyl which may be unsubstituted or substituted by one or more substituents as defined below. Examples of other suitable aryl groups include biphenyl, fluorenyl and indenyl.

The term "heteroaryl" refers to an aromatic ring or ring system that contains one or more heteroatoms, in which the heteroatoms are selected from nitrogen, oxygen and sulfur. Suitable heteroaryl groups include furyl, thienyl, pyridyl, quinolinyl, tetrazolyl, imidazo, and so on. In the case where R_3 and R_4 are taken together and form a 5- or 6-membered heteroaromatic ring, the heteroatom is nitrogen, oxygen or sulfur and the ring may contain one or more of such atoms. Preferably, the heteroatom is nitrogen or sulfur. Preferred heteroaromatic rings formed by R_3 and R_4 are illustrated by the following formulae where the two lines indicate where they are fused.



The terms "heterocyclic" and "heterocyclyl" refer to non-aromatic rings or ring systems that contain one or more ring heteroatoms (e.g., O, S, N). Exemplary heterocyclic groups include pyrrolidinyl, tetrahydrofuranyl, morpholinyl, piperidino, piperazino, thiazolidinyl, imidazolidinyl, and the like.

All of the above rings and ring systems can be unsubstituted or substituted by one or more substituents selected from the group consisting of alkyl, alkoxy, alkylthio, hydroxy, halogen, haloalkyl, polyhaloalkyl, perhaloalkyl (e.g., trifluoromethyl), trifluoroalkoxy (e.g., trifluoromethoxy), nitro, amino, alkylamino, dialkylamino, alkylcarbonyl, alkenylcarbonyl, arylcarbonyl, heteroarylcarbonyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocycloalkyl, nitrile and alkoxycarbonyl. Preferred substituents are C₁₋₄ alkyl, C₁₋₄ alkoxy, halo, amino, alkylamino, dialkylamino, hydroxy, C₁₋₄ alkoxymethyl and trifluoromethyl.

The term "halo" refers to a halogen atom, such as, for example, fluorine, chlorine, bromine or iodine.

The invention is inclusive of the compounds described herein in any of their pharmaceutically acceptable forms, including isomers such as diastereomers and enantiomers, salts, solvates, polymorphs and the like.

As noted above, the compounds of Formula I and I(a) are capable of forming "pharmaceutically acceptable salt(s)." Pharmaceutically acceptable acid addition salts of the compounds of Formula I and I(a) include salts derived from nontoxic inorganic acids such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydriodic, hydrofluoric, phosphorous, and the like, as well as the salts derived from nontoxic organic acids, such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc.



Such salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, trifluoroacetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, hydroxynaphthoate, xinafoate, dinitrobenzoate, phthalate, benzenesulfonate, toluenesulfonate, phenylacetate, citrate, lactate, maleate, tartrate, methanesulfonate, and the like. Also contemplated are salts of amino acids such as arginate and the like and gluconate, galacturonate (see, for example, Berge SM, et al., "Pharmaceutical Salts," *J. Pharm. Sci.* 1977;66:1).

The acid addition salts of the compounds are prepared by contacting the free base form with a sufficient amount of the desired acid to produce the salt in the conventional manner. The free base form may be regenerated by contacting the salt form with a base and isolating the free base in the conventional manner.

Preferred compounds of Formula I and I(a) are those wherein R_1 is oxygen or sulfur. Preferred R_2 substituents include alkyl and alkoxyalkyl, with C_{1-4} alkyl especially preferred.

It is preferred that R_3 and R_4 be taken together to form a fused benzene or pyridine ring that may be substituted or unsubstituted.

Most preferred compounds are those of the Formula III or IV



wherein R₂ is defined above, and R is hydrogen, alkyl, alkoxy, alkylthio, hydroxy, halogen, haloalkyl, polyhaloalkyl, perhaloalkyl (e.g., trifluoromethyl), trifluoroalkoxy (e.g., trifluoromethoxy), nitro, amino, alkylamino, dialkylamino, alkylcarbonyl, alkenylcarbonyl, arylcarbonyl, heteroarylcarbonyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, heterocyclyl, heterocycloalkyl, nitrile and alkoxycarbonyl.

Exemplary compounds of the invention include:

- 2-methylthiazolo[4,5-c]quinolin-4-amine;
- thiazolo[4,5-c]quinolin-4-amine;
- 2-ethylthiazolo[4,5-c]quinolin-4-amine;
- 2-propylthiazolo[4,5-c]quinolin-4-amine;
- 2-pentylthiazolo[4,5-c]quinolin-4-amine;
- 2-butylthiazolo[4,5-c]quinolin-4-amine;
- 2-(1-methylethyl)thiazolo[4,5-c]quinolin-4-amine;
- 2-(2-phenyl-1-ethenyl)thiazolo[4,5-c]quinolin-4-amine;
- 2-(2-phenyl-1-ethyl)thiazolo[4,5-c]quinolin-4-amine;
- 2-(4-aminothiazolo[4,5-c]quinolin-2-yl)-1,1-dimethylethyl carbamate;
- 2-(ethoxymethyl)thiazolo[4,5-c]quinolin-4-amine;
- 2-(methoxymethyl)thiazolo[4,5-c]quinolin-4-amine;
- 2-(2-methylpropyl)thiazolo[4,5-c]quinolin-4-amine;
- 2-benzylthiazolo[4,5-c]quinolin-4-amine;
- 8-methyl-2-propylthiazolo[4,5-c]quinolin-4-amine;
- (4-aminothiazolo[4,5-c]quinolin-2-yl)methanol;
- 2-methyloxazolo[4,5-c]quinolin-4-amine;
- 2-ethyloxazolo[4,5-c]quinolin-4-amine;
- 2-butyloxazolo[4,5-c]quinolin-4-amine;
- 2-propylthiazolo [4,5-c] quinolin-4,8-diamine;
- 2-propyloxazolo [4,5-c] quinolin-4-amine;
- 8-bromo-2-propylthiazolo[4,5-c]quinolin-4-amine;
- 7-methyl-2-propylthiazolo[4,5-c]quinolin-4-amine;
- 2-butyl-7-methyloxazolo[4,5-c]quinolin-4-amine;

7-methyl-2-propyloxazolo[4,5-c]quinolin-4-amine;
7-fluoro-2-propyloxazolo[4,5-c]quinolin-4-amine;
7-fluoro-2-propylthiazolo[4,5-c]quinolin-4-amine;
2-propyl-7-(trifluoromethyl)thiazolo[4,5-c]quinolin-4-amine;
5 2-(4-morpholino)thiazolo[4,5-c]quinolin-4-amine;
2-(1-pyrrolidino)thiazolo[4,5-c]quinolin-4-amine;
2-butylthiazolo[4,5-c][1,5]naphthyridin-4-amine;
2-propylthiazolo[4,5-c][1,5]naphthyridin-4-amine;
7-chloro-2-propylthiazolo[4,5-c]quinolin-4-amine;
10 7-methoxy-2-propylthiazolo[4,5-c]quinolin-4-amine;
and pharmaceutically acceptable salts thereof, particularly the hydrochloride salts thereof.

Preparation of the Compounds

15 Compounds of the invention can be prepared according to Reaction Scheme I wherein R_1 , R_2 , R_3 and R_4 are as defined above.

In step (1) of Reaction Scheme I a compound of Formula V is reacted with a carboxylic acid or an equivalent thereof to provide a compound of Formula VI. Suitable
20 equivalents to carboxylic acid include acid anhydrides, acid chlorides, orthoesters and 1,1-dialkoxyalkanoates. The carboxylic acid or equivalent is selected such that it will provide the desired R_2 substituent in a compound of Formula VI. For example, triethyl orthoformate will provide a compound of Formula VI where R_2 is hydrogen and acetic anhydride will provide a compound of Formula VI where R_2 is methyl. The reaction can
25 be run in the absence of solvent, in the presence of an acid such as polyphosphoric acid, or preferably in the presence of a carboxylic acid of the formula $R_2C(O)OH$. The reaction is run with sufficient heating to drive off any alcohol or water formed as a byproduct of the reaction. The compounds of Formula V are known or may be prepared using conventional methods (see for example, Bachman et. al., Journal of the American Chemical Society, 69,
30 pp 365-371 (1947); Ambrogi et. al., Synthesis, pp. 656-658 (1992); Adler et. al., Journal of the Chemical Society, pp.1794-1797 (1960); Süss et. al., Justus Liebigs Annalen der

Chemie, 583, pp. 150-160 (1953); and Süss et. al., *Justus Liebigs Annalen der Chemie*, 593, pp. 91-126 (1955).


In step (2) of Reaction Scheme I a compound of Formula VI is oxidized to provide an N-oxide of Formula II. The oxidation is carried out using a conventional oxidizing agent that is capable of forming N-oxides. Preferred reaction conditions involve reacting a solution of a compound of Formula VI in chloroform with 3-chloroperoxybenzoic acid at ambient conditions. Alternatively the oxidation may be carried out using peracetic acid in a suitable solvent such as ethyl or methyl acetate.

In step (3) of Reaction Scheme I an N-oxide of Formula II is aminated to provide a compound of Formula I. Step (3) involves (i) reacting a compound of Formula II with an acylating agent and then (ii) reacting the product with an aminating agent. Part (i) of step (3) involves reacting an N-oxide of Formula II with an acylating agent. Suitable acylating agents include alkyl- or arylsulfonyl chlorides (e.g., benzenesulfonyl chloride, methanesulfonyl chloride, p-toluenesulfonyl chloride). Arylsulfonyl chlorides are preferred. *Para*-toluenesulfonyl chloride is most preferred. Part (ii) of step (3) involves reacting the product of part (i) with an excess of an aminating agent. Suitable aminating agents include ammonia (e.g., in the form of ammonium hydroxide) and ammonium salts (e.g., ammonium carbonate, ammonium bicarbonate, ammonium phosphate). Ammonium hydroxide is preferred. The reaction is preferably carried out by dissolving or suspending the N-oxide of Formula II in an inert solvent such as dichloromethane or chloroform, adding the aminating agent to the solution or suspension, and then slowly adding the acylating agent. The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Alternatively, step (3) can be carried out by (i) reacting an N-oxide of Formula II with an isocyanate and then (ii) hydrolyzing the resulting product. Part (i) involves reacting the N-oxide with an isocyanate wherein the isocyanato group is bonded to a carbonyl group. Preferred isocyanates include trichloroacetyl isocyanate and aroyl isocyanates such as benzoyl isocyanate. The reaction of the isocyanate with the N-oxide is carried out under substantially anhydrous conditions by adding the isocyanate to a solution of the N-oxide in an inert solvent such as dichloromethane. Part (ii) involves hydrolysis of the product from part (i). The hydrolysis can be carried out by conventional methods such

as heating in the presence of water or a lower alkanol optionally in the presence of a catalyst such as an alkali metal lower alkoxide or ammonia.

Reaction Scheme I



5

Compounds of the invention wherein R_1 is oxygen or sulfur and R_3 and R_4 together form an optionally substituted aromatic ring can be prepared according to Reaction Scheme II wherein R and R_2 are as defined above.

10

In step (1) of Reaction Scheme II a 3-aminoquinolin-4-ol or 3-aminoquinolin-4-thiol of Formula VII is reacted with a carboxylic acid or an equivalent thereof to provide an oxazolo- or thiazolo[4,5-*c*]quinoline of Formula VIII. Suitable equivalents to carboxylic acid include acid anhydrides, acid chlorides, orthoesters and 1,1-dialkoxyalkanoates. The carboxylic acid or equivalent is selected such that it will provide the desired R_2 substituent in a compound of Formula VIII. For example, triethyl orthoformate will provide a compound of Formula VIII where R_2 is hydrogen and acetic anhydride will provide a compound of Formula VIII where R_2 is methyl. The reaction can be run in the absence of solvent, in the presence of an acid such as polyphosphoric acid, or preferably in the presence of a carboxylic acid of the formula $R_2C(O)OH$. The reaction is run with sufficient heating to drive off any alcohol or water formed as a byproduct of the

20

reaction. The 3-aminoquinolin-4-ols and 3-aminoquinolin-4-thiols of Formula VII are known or may be prepared using known methods.

In step (2) of Reaction Scheme II an oxazolo- or thiazolo[4,5-*c*]quinoline of Formula VIII is oxidized to provide an oxazolo- or thiazolo[4,5-*c*]quinolin-5N-oxide of Formula IX which is a subgenus of Formula II. The oxidation is carried out using a conventional oxidizing agent that is capable of forming N-oxides. Preferred reaction conditions involve reacting a solution of a compound of Formula VIII in chloroform with 3-chloroperoxybenzoic acid at ambient conditions. Alternatively the oxidation may be carried out using peracetic acid in a suitable solvent such as ethyl or methyl acetate.


In step (3) of Reaction Scheme II an N-oxide of Formula IX is aminated to provide an oxazolo[4,5-*c*]quinolin-4-amine of Formula III or a thiazolo[4,5-*c*]quinolin-4-amine of Formula IV both of which are subgenera of Formula I. Step (3) involves (i) reacting a compound of Formula IX with an acylating agent and then (ii) reacting the product with an aminating agent. Part (i) of step (3) involves reacting an N-oxide of Formula IX with an acylating agent. Suitable acylating agents include alkyl- or arylsulfonyl chlorides (e.g., benzenesulfonyl chloride, methanesulfonyl chloride, *p*-toluenesulfonyl chloride). Arylsulfonyl chlorides are preferred. *Para*-toluenesulfonyl chloride is most preferred. Part (ii) of step (3) involves reacting the product of part (i) with an excess of an aminating agent. Suitable aminating agents include ammonia (e.g., in the form of ammonium hydroxide) and ammonium salts (e.g., ammonium carbonate, ammonium bicarbonate, ammonium phosphate). Ammonium hydroxide is preferred. The reaction is preferably carried out by dissolving or suspending the N-oxide of Formula IX in an inert solvent such as dichloromethane or chloroform, adding the aminating agent to the solution or suspension, and then slowly adding the acylating agent. The product or a pharmaceutically acceptable salt thereof can be isolated using conventional methods.

Alternatively, step (3) can be carried out by (i) reacting an N-oxide of Formula IX with an isocyanate and then (ii) hydrolyzing the resulting product. Part (i) involves reacting the N-oxide with an isocyanate wherein the isocyanato group is bonded to a carbonyl group. Preferred isocyanates include trichloroacetyl isocyanate and aroyl isocyanates such as benzoyl isocyanate. The reaction of the isocyanate with the N-oxide is carried out under substantially anhydrous conditions by adding the isocyanate to a solution

of the N-oxide in an inert solvent such as dichloromethane. Part (ii) involves hydrolysis of the product from part (i). The hydrolysis can be carried out by conventional methods such as heating in the presence of water or a lower alkanol optionally in the presence of a catalyst such as an alkali metal lower alkoxide or ammonia.

5

Reaction Scheme II



Compounds of the invention wherein R_1 is sulfur can also be prepared according to Reaction Scheme III wherein R_2 , R_3 and R_4 are as defined above.

10


In step (1) of Reaction Scheme III a compound of Formula X is reacted with an acyl halide of formula $\text{R}_2\text{C}(\text{O})\text{Z}$ wherein R_2 is as defined above and Z is chloro or bromo to provide an amide of Formula XI. The reaction can be carried out by adding the acyl halide in a controlled fashion (e.g., dropwise) to a solution or suspension of a compound of Formula X in a suitable solvent such as pyridine or dichloromethane in the presence of a tertiary amine.

15

In step (2) of Reaction Scheme III an amide of Formula XI is reacted with phosphorous pentasulfide to provide a compound of Formula XII. The reaction can be

carried out by adding phosphorous pentasulfide to a solution or suspension of a compound of Formula XI in a suitable solvent such as pyridine and heating the resulting mixture.

Steps (3) and (4) of Reaction Scheme III can be carried out in the same manner as steps (2) and (3) of Reaction Scheme I respectively to provide an N-oxide of Formula XIII which is a subgenus of Formula II and a compound of Formula XIV which is a subgenus of Formula I respectively.




Compounds of the invention wherein R_1 is sulfur and R_3 and R_4 together form an optionally substituted aromatic ring can also be prepared according to Reaction Scheme IV wherein R and R_2 are as defined above.


In step (1) of Reaction Scheme IV a 3-aminoquinolin-4-ol of Formula XV is reacted with an acyl halide of formula $R_2C(O)Z$ wherein R_2 is as defined above and Z is chloro or bromo to provide an N-(4-hydroxyquinolin-3-yl)amide of Formula XVI. The reaction can be carried out by adding the acyl halide in a controlled fashion (e.g., dropwise) to a solution or suspension of a compound of Formula XV in a suitable solvent such as dichloromethane in the presence of a tertiary amine.

In step (2) of Reaction Scheme IV an N-(4-hydroxyquinolin-3-yl)amide of Formula XVI is reacted with phosphorous pentasulfide to provide a thiazolo[4,5-c]quinoline of Formula XVII. The reaction can be carried out by adding phosphorous pentasulfide to a solution or suspension of a compound of Formula XVI in a suitable solvent such as pyridine and heating the resulting mixture.

Steps (3) and (4) of Reaction Scheme IV can be carried out in the same manner as steps (2) and (3) of Reaction Scheme II respectively to provide a thiazolo[4,5-c]quinolin-5N-oxide of Formula XVIII which is a subgenus of Formula II and a thiazolo[4,5-c]quinolin-4-amine of Formula IV which is a subgenus of Formula I respectively.



Substituents at the 2-position can be introduced by reacting a compound of Formula XIX



wherein R₁ is oxygen or sulfur and R is as defined above, with a lithiating agent such as lithium diisopropylamide or n-butyllithium in a polar aprotic solvent to provide a compound lithiated on the 2-methyl group. The lithiated compound can then be reacted with an appropriate reagent containing a leaving group capable of being displaced by the lithiated 2-methyl group. Examples of suitable reagents include halides such as methyl iodide or chloromethylmethylether, aldehydes such as benzaldehyde, and ketones such as acetone. The compounds can then be oxidized and aminated using the methods described above to provide compounds of Formulas III or IV.

Some compounds of Formula I may be prepared directly from other compounds of Formula I. For example, nitration of 2-propylthiazolo[4,5-c]quinolin-4-amine provides 8-nitro-2-propylthiazolo[4,5-c]quinolin-4-amine and the reduction of the nitro compound provides 2-propylthiazolo[4,5-c]quinoline-4,8-diamine.

Pharmaceutical Compositions and Biological Activity

Pharmaceutical compositions of the invention contain a therapeutically effective amount of a compound of Formula I(a) together with a pharmaceutically acceptable carrier.

As used herein, the term "a therapeutically effective amount" means an amount of the compound sufficient to induce a desired therapeutic effect, such as cytokine biosynthesis, antitumor activity and/or antiviral activity. Although the exact amount of active compound used in a pharmaceutical composition of the invention will vary according to factors known to those of skill in the art, such as the physical and chemical nature of the compound as well as the nature of the carrier, the intended dosing regimen and the condition to be treated, it is anticipated that the compositions of the invention will contain sufficient active ingredient to provide a dose of about 100ng/kg to about 50mg/kg, preferably about 10μg/kg to about 5mg/kg of the compound to the subject. Any of the conventional dosage forms may be used, such as tablets, lozenges, parenteral formulations, syrups, creams, ointments, aerosol formulations, transdermal patches, transmucosal patches and so on. The dosage form used will also depend on the characteristics of the compound to be administered. For example, certain compounds of Formula I(a), especially those wherein R₁ is sulfur, tend to have relatively low oral bioavailability and

are rapidly metabolized when they enter the bloodstream. These properties make such compounds particularly well suited for treatment of conditions where topical administration of an immune response modifying compound is desirable, such as asthma, basal cell carcinoma, cervical intraepithelial neoplasia and so on.

5 The compounds of the invention have been shown to induce the production of certain cytokines in experiments performed according to the Test Method set forth below. These results indicate that the compounds are useful as immune response modifiers that can modulate the immune response in a number of different ways, rendering them useful in the treatment of a variety of disorders.

10 Cytokines that are induced by the administration of compounds according to the invention generally include interferon- α (IFN- α) and/or tumor necrosis factor- α (TNF- α) as well as certain interleukins (IL). Cytokines whose biosynthesis may be induced by compounds of the invention include IFN- α , TNF- α , IL-1, 6, 10 and 12, and a variety of other cytokines. Among other effects, cytokines inhibit virus production and tumor cell
15 growth, making the compounds useful in the treatment of tumors and viral diseases.

 In addition to the ability to induce the production of cytokines, the compounds of the invention affect other aspects of the innate immune response. For example, natural killer cell activity may be stimulated, which effect may be due to cytokine induction. The compounds may also activate macrophages, which in turn stimulates secretion of nitric
20 oxide and the production of additional cytokines. Further, the compounds may cause proliferation and differentiation of B-lymphocytes and may be useful in the *in vitro* maturation of dendritic cells.

 Compounds of the invention also have an effect on the acquired immune response. For example, although there is not believed to be any direct effect on T cells or direct
25 induction of T cell cytokines, the production of the T helper type 1 (Th1) cytokine IFN- γ is induced indirectly and the production of the T helper type 2 (Th2) cytokines IL-4, IL-5 and IL-13 are inhibited upon administration of the compounds. This activity means that the compounds are useful in the treatment of diseases where upregulation of the Th1 response and/or down regulation of the Th2 response is desired. In view of the ability of
30 compounds of Formula Ia to inhibit the Th2 immune response, the compounds are expected to be useful in the treatment of atopy, e.g., atopic dermatitis, asthma, allergy,

allergic rhinitis, and systemic lupus erythematosus; as a vaccine adjuvant for the enhancement of cell mediated immunity; and possibly as a treatment for recurrent fungal diseases and chlamydia.

The immune response modifying effects of the compounds make them useful in the treatment of a wide variety of conditions. Because of their ability to induce cytokines such as IFN- α and/or TNF- α , the compounds are particularly useful in the treatment of viral diseases and tumors. This immunomodulating activity suggests that compounds of the invention are useful in treating diseases such as, but not limited to, viral diseases e.g., genital warts, common warts, plantar warts, Hepatitis B, Hepatitis C, Herpes Simplex Type I and Type II, molluscum contagiosum, HIV, CMV, VZV, cervical intraepithelial neoplasia, human papillomavirus and associated neoplasias; fungal diseases, e.g., candida, aspergillus, cryptococcal meningitis; neoplastic diseases, e.g., basal cell carcinoma, hairy cell leukemia, Kaposi's sarcoma, renal cell carcinoma, squamous cell carcinoma, myelogenous leukemia, multiple myeloma, melanoma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, and other cancers; parasitic diseases, e.g., pneumocystis carinii, cryptosporidiosis, histoplasmosis, toxoplasmosis, trypanosome infection, leishmaniasis; and bacterial infections, e.g., tuberculosis, mycobacterium avium. Additional diseases or conditions that can be treated using the compounds of the invention include eczema, eosinophilia, essential thrombocythaemia, leprosy, multiple sclerosis, Ommen's syndrome, rheumatoid arthritis, systemic lupus erythematosus, discoid lupus, Bowen's disease, Bowenoid papulosis, and to enhance or stimulate the healing of wounds, including chronic wounds.

Accordingly, the invention provides a method of inducing cytokine biosynthesis in an animal comprising administering an effective amount of a compound of Formula Ia to the animal. An amount of a compound effective to induce cytokine biosynthesis is an amount sufficient to cause one or more cell types, such as monocytes, macrophages, dendritic cells and B-cells to produce an amount of one or more cytokines such as, for example, IFN- α , TNF- α , IL-1,6,10 and 12 that is increased over the background level of such cytokines. The precise amount will vary according to factors known in the art but is expected to be a dose of about 100 ng/kg to about 50mg/kg, preferably about 10 μ g/kg to about 5mg/kg. The invention also provides a method of treating a viral infection in an

animal comprising administering an effective amount of a compound of Formula Ia to the animal. An amount effective to treat or inhibit a viral infection is an amount that will cause a reduction in one or more of the manifestations of viral infection, such as viral lesions, viral load, rate of virus production, and mortality as compared to untreated control animals. The precise amount will vary according to factors known in the art but is expected to be a dose of 100ng/kg to about 50mg/kg, preferably about 10µg/kg to about 5mg/kg.


Compounds of the invention may be administered to the subject as the sole therapeutic agent, or may form part of a therapeutic regimen in combination with one or more other agents. Examples of suitable agents that may be used in combination with the immune response modifying compounds of the invention include, but are not limited to, analgesics, antibacterials, antifungals, antiinflammatory agents, antitumor agents, antivirals, bronchodilators, narcotics, and steroids.

The following examples are provided to illustrate the invention, but should not be considered to be limiting in any way.

EXAMPLES

Example 1

2-Methylthiazolo[4,5-c]quinoline-5N-oxide



Part A

A suspension of 3-aminoquinoline-4-thiol (about 12 g) in a mixture of acetic anhydride (150 mL) and acetic acid (300 mL) was heated at reflux overnight. The reaction mixture was filtered to remove a fine solid. The filtrate was evaporated under vacuum. The residue was diluted with ethanol then refluxed for 30 minutes. The solution was concentrated under vacuum and the residue diluted with water. The aqueous residue was made basic with sodium hydroxide and then extracted with diethyl ether. The ether extracts were combined, dried over magnesium sulfate and filtered. The filtrate was


evaporated to provide 12.8 g of crude product. A sample (800 mg) was recrystallized from hexane to provide 2-methylthiazolo[4,5-*c*]quinoline as yellow needles, m.p. 95.5-97.5°C.^a Analysis: Calculated for C₁₁H₈N₂S: %C, 65.97; %H, 4.03; %N, 13.99; Found: %C, 65.96; %H, 4.16; %N 14.08.

5 Part B

2-Methylthiazolo[4,5-*c*]quinoline (5.0 g, 25 mmol), 3-chloroperoxybenzoic acid (9.5 g of 50-60%), and dichloromethane (150 mL) were combined and stirred at ambient temperature for 3 hours. The reaction solution was diluted with dichloromethane (300 mL) and then extracted with aqueous sodium carbonate to remove the acids. The organic
10 layer was washed with water, diluted with ethyl acetate to remove cloudiness, dried over magnesium sulfate, and then concentrated under vacuum to provide 4.5 g of crude product. A small portion was recrystallized from methanol to provide yellow needles of 2-methylthiazolo[4,5-*c*]quinoline-5N-oxide hydrate, m.p. 150-160°C. Analysis: Calculated for C₁₁H₁₈N₂OS + 0.75 H₂O: %C, 57.50; %H, 4.17; %N, 12.19; Found: %C, 57.58; %H,
15 4.10; %N, 11.93.

Example 2

2-Methylthiazolo[4,5-*c*]quinolin-4-amine



20

2-Methylthiazolo[4,5-*c*]quinoline-5N-oxide (1.5 g, 6.9 mmol) was added to a mixture of dichloromethane (10 mL) and ammonium hydroxide (25 mL). A solution of tosyl chloride (2.0 g, 10.4 mmol) in dichloromethane (10 mL) was added with vigorous stirring to the reaction mixture. The reaction was refluxed with additional
25 dichloromethane and ammonium hydroxide being added until thin layer chromatography indicated that the reaction was complete. The dichloromethane was distilled from the mixture and yellow product was filtered from the aqueous residue. The solid was washed

with water and then dried to provide 1.2 g of crude product. The solid was dissolved in dilute hydrochloric acid. The solution was treated with charcoal and then filtered. The filtrate was made basic with dilute sodium hydroxide. The resulting precipitate was isolated by filtration, washed with water, dried and then recrystallized from methanol/dichloromethane to provide 0.46 g of 2-methylthiazolo[4,5-*c*]quinolin-4-amine as a white powder, m.p. 184-187°C. Analysis: Calculated for C₁₁H₉N₃S: %C, 61.37; %H, 4.21; %N, 19.52; Found: %C, 61.32; %H, 4.52; %N, 19.68.

Example 3

2-Methylthiazolo[4,5-*c*]quinolin-4-amine

Alternative Synthesis

Trichloroacetyl isocyanate (2.0 mL, 16.8 mmol) was added to a suspension of 2-methylthiazolo[4,5-*c*]quinoline-5N-oxide (3.03 g, 14.0 mmol) in dichloromethane (150 mL). The reaction mixture was stirred at ambient temperature for about 50 minutes. The dichloromethane was concentrated under vacuum to provide crude N-(2-methylthiazolo[4,5-*c*]quinolin-4-yl)trichloroacetamide. The amide was dissolved in methanol and then sodium methoxide (1 mL of 25% sodium methoxide in methanol) was added. The reaction was heated at reflux for 40 minutes then the methanol was evaporated under vacuum. The resulting brown solid was washed with water and dried to provide 2.85 g of crude product. This material was treated with charcoal and then recrystallized from ethyl acetate to provide 2-methylthiazolo[4,5-*c*]quinolin-4-amine as a solid, m.p. 184-186°C. Analysis: Calculated for C₁₁H₉N₃S: %C, 61.37; %H, 4.21; %N, 19.52; Found: %C, 61.48; %H, 4.17; %N, 19.60.

Example 4

2-Methylthiazolo[4,5-*c*]quinolin-4-amine Hydrochloride

Concentrated hydrochloric acid (0.2 mL of 12.1M) was added to a solution of 2-methylthiazolo[4,5-*c*]quinolin-4-amine (0.5 g) in methanol (15 mL). Isopropanol (15 mL) was added and then the reaction mixture was heated at reflux to remove the majority of the methanol. The resulting precipitate was isolated by filtration, washed with isopropanol and dried to provide 2-methylthiazolo[4,5-*c*]quinolin-4-amine hydrochloride as a solid,

m.p. 323-325°C. Analysis: Calculated for $C_{11}H_9N_3S \cdot HCl$: %C, 52.48; %H, 4.00; %N, 16.69; Found: %C, 52.46; %H, 4.08; %N, 16.52.

Example 5

5 Thiazolo[4,5-*c*]quinolin-4-amine Hydrochloride Hydrate

Part A

3-Aminoquinoline-2-thiol (about 18.5 g) was added to triethyl orthoformate (26.0 mL). The reaction mixture was heated on a steam bath for 20 minutes. Formic acid (400 mL) was added and the reaction mixture was heated at reflux overnight. The bulk of the
10 formic acid was evaporated under vacuum. The residue was combined with ethanol and heated at reflux for 30 minutes. The ethanol was evaporated under vacuum. The residue was suspended in water and then made basic by adding sodium hydroxide. A precipitate formed. The solid was extracted with several portions of dichloromethane. The extracts were combined, dried over magnesium sulfate and then concentrated to provide a yellow
15 solid which was recrystallized from hexanes to provide 13.1 g of thiazolo[4,5-*c*]quinoline as a yellow crystalline solid, m.p. 104-106°C.

Part B

Peracetic acid (21 mL of 32% in acetic acid, 100 mmol) was added to a suspension of thiazolo[4,5-*c*]quinoline (12.5 g, 67 mmol) in methyl acetate (300 mL). The reaction
20 mixture was heated at reflux overnight and then cooled to ambient temperature. A precipitate was isolated by filtration and then suspended in water (100 mL). Aqueous sodium bicarbonate (100 mL) was added to the suspension and the mixture was stirred for one hour. The solid thiazolo[4,5-*c*]quinoline-5N-oxide was isolated by filtration, washed with water and dried.


25 Part C

Trichloroacetyl isocyanate (0.72 mL, 6.0 mmol) was added to a suspension of thiazolo[4,5-*c*]quinoline-5N-oxide (1.10 g, 5.4 mmol) in dichloromethane (100 mL). The reaction mixture was stirred at ambient temperature for 45 minutes and then the dichloromethane was evaporated under vacuum to provide crude N-(thiazolo[4,5-
30 *c*]quinolin-4-yl)trichloroacetamide. The amide was combined with 2M ammonia in methanol and stirred at ambient temperature for 2 hours. The methanol was evaporated

under vacuum. The residue was suspended in water, combined with sodium carbonate and then stirred for 10 minutes. A brown solid was collected, washed with water and dried.[•] The solid was suspended in water, hydrochloric acid (100 mL of 6N) was added and the mixture was heated on a steam bath. The mixture was filtered; then the filtrate was
5 allowed to slowly cool to ambient temperature. The resulting precipitate was isolated by filtration and then dried to provide 0.75 g of brown needles. This material was dissolved in water (100 mL) with heating. Charcoal was added and the mixture was heated with stirring for 5 minutes. The mixture was filtered through a layer of Celite® filter agent. The filtrate was heated on a steam bath to remove most of the water and then allowed to
10 cool to ambient temperature. The precipitate was isolated by filtration and dried to provide 0.30 g of thiazolo[4,5-*c*]quinolin-4-amine hydrochloride hydrate as a white crystalline solid, m.p. 284-285°C. Analysis: Calculated for C₁₀H₇N₃S · HCl · H₂O: %C, 46.97; %H, 3.94; %N, 16.43; Found: %C, 46.96; %H, 3.99; %N, 16.34.

15

Example 6

Thiazolo[4,5-*c*]quinolin-4-aminePart A

20

Thiazolo[4,5-*c*]quinoline-2-thiol (8.7 g, 0.04 mole) was suspended in a solution of sodium hydroxide (1.4 g, 0.04 mole) in water. A few drops of 50% sodium hydroxide were added to the suspension until most of the solid dissolved. Hydrogen peroxide (13.5 mL of 30%, 0.08 mole) was added dropwise over a period of 30 minutes while maintaining the temperature of the reaction mixture at 25-35°C with a cold water bath.

25

The bath was removed and the reaction mixture was stirred for 15 minutes. Sulfuric acid (2.5 g of 95.98%) was added dropwise to the reaction mixture. After 30 minutes the reaction was made basic (pH 9-9.5) with 50% sodium hydroxide. The reaction mixture

was acidified (pH 2.5) with hydrochloric acid, a tan solid precipitated. The mixture was then heated on a steam bath for 15 minutes and the precipitate dissolved. The solution was allowed to cool to ambient temperature and a precipitate formed. The mixture was made basic (pH 9) with 50% sodium hydroxide. The resulting oily product was extracted with ethyl acetate. The extracts were combined, washed with water, dried over magnesium sulfate and then concentrated under vacuum to provide 3.3 g of thiazolo[4,5-*c*]quinoline as a tan solid, m.p. 104.4-105°C. Analysis: Calculated for C₁₀H₆N₂S: %C, 64.19; %H, 3.25; %N, 15.04; Found: %C, 64.15; %H, 3.26; %N, 14.9.

Part B

Peracetic acid (4.7 mL of 32%) was added to a solution of thiazolo[4,5-*c*]quinoline (2.8 g) in methyl acetate. A precipitate formed after several minutes. The reaction mixture was heated to reflux and then diluted with an additional 10 mL of methyl acetate. Most of the precipitate dissolved. After 1 hour an additional 3.1 mL of peracetic acid was added. The reaction mixture was heated overnight and then allowed to cool to ambient temperature. The methyl acetate and acetic acid were azeotroped off with heptane. The resulting oily product was suspended in water. The mixture was made basic with saturated sodium bicarbonate and then extracted with ethyl acetate. The ethyl acetate extracts were combined, washed with water, dried over magnesium sulfate and then concentrated under vacuum to provide 0.6 g of thiazolo[4,5-*c*]quinoline-5N-oxide as an orange solid, m.p. 178.4°C (dec).

Part C


A solution of tosyl chloride (0.3 g) in water was added dropwise to a cooled (5°C) suspension of thiazolo[4,5-*c*]quinoline-5N-oxide (0.3 g) in a mixture of ammonium hydroxide (5 mL) and dichloromethane (50 mL). The temperature was maintained at 4-6°C throughout the addition. After the addition was complete the reaction mixture was stirred at ambient temperature for 4 hours. Analysis by thin layer chromatography showed the presence of starting material. The reaction mixture was cooled and 1 equivalent of tosyl chloride was added. The reaction mixture was stirred at ambient temperature for 20 hours. The dichloromethane was evaporated under vacuum. The residue was slurried in a small amount of water. The mixture was filtered. The isolated solid was washed with water, dried and then recrystallized from isopropanol to provide 0.2 g of thiazolo[4,5-

c]quinolin-4-amine as an orange powder, m.p. 172.4°C (dec). Analysis: Calculated for $C_{10}H_7N_3S$: %C, 59.68; %H, 3.50; %N, 20.88; Found: %C, 59.82; %H, 3.20; %N, 19.50.

Example 7

5

2-Ethylthiazolo[4,5-c]quinoline-5N-oxide



Part A

Under a nitrogen atmosphere, 2-methylthiazolo[4,5-c]quinoline (1.0 g, 10.0 mmol, Example 2 or 3) was placed in a dried flask. Anhydrous tetrahydrofuran (50 mL) was added and the reaction mixture was cooled to -78°C with a dry ice bath. Lithium diisopropylamide (6.7 mL of 1.5 M in hexane, 10.0 mmol) was added dropwise. 30 minutes later methyl iodide (0.95 mL, 15.0 mmol) was added. After 40 minutes the reaction was allowed to warm to ambient temperature. The reaction mixture was quenched with water and then extracted with diethyl ether (250 mL). The extract was washed with water (3 X 100 mL), dried over magnesium sulfate and then concentrated under vacuum to provide 2.8 g of a brown oil. This material was purified using high performance liquid chromatography eluting with 3:1 hexane:ethyl acetate to provide 1.47 g of 2-ethylthiazolo[4,5-c]quinoline as a yellow oil.


20

Part B

3-Chloroperoxybenzoic acid (0.44 g) was added to a solution of 2-ethylthiazolo[4,5-c]quinoline (0.53 g) in chloroform (20 mL). The reaction mixture was stirred at ambient temperature for 2 hours and then diluted with dichloromethane (20 mL), washed with sodium bicarbonate, washed with water (3 X 100 mL), dried over magnesium sulfate and then concentrated under vacuum. The resulting yellow solid was recrystallized from ethyl acetate to provide 0.32 g of 2-ethylthiazolo[4,5-c]quinoline-5N-oxide as a solid, m.p. 128°C. Analysis: Calculated for $C_{12}H_{10}N_2OS$: %C, 62.59 %H, 4.38; %N, 12.16; Found: %C, 62.59; %H, 4.27; %N, 12.12.

Example 8

2-Ethylthiazolo[4,5-c]quinolin-4-amine



5

Part A

Trichloroacetyl isocyanate (0.51 mL, 4.3 mmol) was added to a suspension of 2-ethylthiazolo[4,5-c]quinoline-5N-oxide (0.90 g, 3.9 mmol) in dichloromethane (60 mL). The reaction mixture was stirred at ambient temperature for 30 minutes and then concentrated under vacuum to provide 1.80 g of N-(2-ethylthiazolo[4,5-c]quinolin-4-yl)trichloroacetamide as a yellow solid.

10

Part B

N-(2-ethylthiazolo[4,5-c]quinolin-4-yl)trichloroacetamide (0.40 g) was suspended in a solution of ammonia in methanol (20 mL of 2M) and then stirred at ambient temperature for 30 minutes. The reaction mixture was concentrated under vacuum. The residue was washed with water and then dried to provide 0.19 g of crude product which was recrystallized from ethyl acetate/hexane to provide 2-ethylthiazolo[4,5-c]quinolin-4-amine as tan needles, m.p. 170-172°C. Analysis: Calculated for C₁₂H₁₁N₃S: %C, 62.85; %H, 4.83; %N, 18.32; Found: %C, 62.58; %H, 4.78; %N, 18.08.

15

20

Example 9

2-Ethylthiazolo[4,5-c]quinolin-4-amine

Alternative Synthesis

Part A

Propionyl anhydride (20 mL) was added to a suspension of 3-amino-quinoline-4-thiol (15 g) in propionic acid (100 mL). The reaction mixture was heated at reflux overnight and then filtered to remove a precipitate. The filtrate was concentrated under

25

vacuum. The residue was taken up in dichloromethane (200 mL), washed with sodium bicarbonate then with water, and then dried over magnesium sulfate. The solution was filtered through a layer of silica gel eluting first with 1:1 ethyl acetate:hexane and then with ethyl acetate. The filtrate was evaporated to provide 2.6 g of 2-ethylthiazolo[4,5-
5 c]quinoline as a yellow oil.

Part B

Peracetic acid (7.4 mL of 32%) was added to a solution of 2-ethylthiazolo[4,5-
c]quinoline (5 g) in ethyl acetate (100 mL). The reaction mixture was stirred at ambient temperature for 2 days. The resulting precipitate was isolated by filtration, washed with
10 hexane and dried to provide 3.4 g of 2-ethylthiazolo[4,5-c]quinoline-5N-oxide.

Part C


Trichloroacetyl isocyanate (6.5 mL, 54 mmol) was added to a suspension of 2-ethylthiazolo[4,5-c]quinoline-5N-oxide (9.0 g, 39.1 mmol) in dichloromethane (500 mL). The reaction mixture was stirred at ambient temperature for 2 hours and then concentrated
15 under vacuum to provide crude N-(2-ethylthiazolo[4,5-c]quinolin-4-yl)trichloroacetamide. This material was added to a solution of ammonia in methanol (500 mL of 2M) and stirred at ambient temperature for about 2 hours. The reaction mixture was concentrated under vacuum. The residue was taken up in dichloromethane, washed with sodium bicarbonate (2 X 150 mL) then with water (3 X 150 mL), dried over magnesium sulfate and then
20 concentrated under vacuum. The residue was recrystallized from 1,2-dichloroethane to provide tan needles. This material was suspended in water; one equivalent of concentrated hydrochloric acid was added and the mixture was heated to dissolve the solids. The solution was treated with charcoal and then filtered. The filtrate was cooled and then made basic with sodium carbonate. The resulting precipitate was isolated by filtration, washed
25 with water and then recrystallized from 1,2-dichloroethane to provide 2-ethylthiazolo[4,5-c]quinolin-4-amine as yellow needles, m.p. 169-171°C. Analysis: Calculated for $C_{12}H_{11}N_3S$: %C, 62.85; %H, 4.83; %N, 18.32; Found: %C, 62.79; %H, 4.86; %N, 18.22.

Example 10

2-Ethylthiazolo[4,5-*c*]quinolin-4-amine Hydrochloride

Concentrated hydrochloric acid (18.5 mmol) was added to a solution of 2-ethylthiazolo[4,5-*c*]quinolin-4-amine (4.25 g) in warm isopropanol. The reaction mixture was heated at reflux to reduce the volume and remove the water. The reaction mixture was cooled to ambient temperature. The precipitate was isolated by filtration and then dried to provide 2-ethylthiazolo[4,5-*c*]quinolin-4-amine hydrochloride as a solid, m.p. 268-270°C. Analysis: Calculated for C₁₂H₁₁N₃S · HCl: %C, 54.23; %H, 4.55; %N, 15.81; Found: %C, 54.25; %H, 4.63; %N, 15.71.

Example 11

2-Propylthiazolo[4,5-*c*]quinoline-5N-oxidePart A


Using the general method of Example 7 Part A, 2-methylthiazolo[4,5-*c*]quinoline (2.50 g, 12.5 mmol) was reacted first with lithium diisopropylamide and then with ethyl iodide to provide 0.28 g of 2-propylthiazolo[4,5-*c*]quinoline as a yellow crystalline solid, m.p. 54°C. Analysis: Calculated for C₁₃H₁₂N₂S: %C, 68.39; %H, 5.30; %N, 12.27; Found: %C, 68.41; %H, 5.19; %N, 12.31.

Part B

Using the general method of Example 7 Part B, 2-propylthiazolo[4,5-*c*]quinoline (1.05 g, 4.6 mmol) was oxidized with 3-chloroperoxybenzoic acid to provide 0.65 g of 2-propylthiazolo[4,5-*c*]quinoline-5N-oxide as a yellow solid, m.p. 123°C. Analysis: Calculated for C₁₃H₁₂N₂OS: %C, 63.91; %H, 4.95; %N, 11.47; Found: %C, 63.53; %H, 4.88; %N, 11.44.

Example 12

2-Propylthiazolo[4,5-c]quinolin-4-amine



Using the general method of Example 8, 2-propylthiazolo[4,5-c]quinoline-5N-oxide (0.63 g) was reacted with trichloroacetyl isocyanate and the resulting amide intermediate was hydrolyzed using a solution of ammonia in methanol to provide 0.22 g of 2-propylthiazolo[4,5-c]quinolin-4-amine as a white crystalline solid, m.p. 140-142°C. Analysis: Calculated for C₁₃H₁₃N₃S: %C, 64.17; %H, 5.38; %N, 17.27; Found: %C, 64.31; %H, 5.39; %N, 17.13.

10

Example 13

2-Propylthiazolo[4,5-c]quinolin-4-amine

Alternative Synthesis

Part A

Using the general method of Example 9 Part A, a suspension of 3-amino-quinoline-4-thiol (15 g) in butyric acid was reacted with butyric anhydride to provide 2-propylthiazolo[4,5-c]quinoline as a yellow oil.

Part B

Using the general method of Example 9 Part B, 2-propylthiazolo[4,5-c]quinoline (46 g) was oxidized with peracetic acid to provide 2-propylthiazolo[4,5-c]quinoline-5N-oxide as a yellow crystalline solid..

Part C

Ammonium hydroxide (50 mL) was added to a solution of 2-propylthiazolo[4,5-c]quinoline-5N-oxide (20 g) in chloroform (500 mL). The reaction mixture was cooled with an ice bath and then a solution of tosyl chloride (16 g) in chloroform was added dropwise. The reaction mixture was refluxed for 2 hours and then diluted with additional chloroform and water. The layers were separated. The organic layer was washed with

25

aqueous sodium bicarbonate, dried over magnesium sulfate and then concentrated under vacuum. The residue was recrystallized from 1,2-dichloroethane to provide 2-propylthiazolo[4,5-*c*]quinolin-4-amine as a tan solid, m.p. 140-142°C. Analysis: Calculated for C₁₃H₁₃N₃S: %C, 64.17; %H, 5.38; %N, 17.27; Found: %C, 64.10; %H, 5.47; %N, 17.29.


Example 14

2-Propylthiazolo[4,5-*c*]quinolin-4-amine Hydrochloride

Using the general method of Example 10, 2-propylthiazolo[4,5-*c*]quinolin-4-amine (1.75 g) was reacted with 1 equivalent of concentrated hydrochloric acid to provide 2-propylthiazolo[4,5-*c*]quinolin-4-amine hydrochloride as an off-white crystalline solid, m.p. 234-237°C. Analysis: Calculated for C₁₃H₁₃N₃S · HCl: %C, 55.81; %H, 5.04; %N, 15.02; Found: %C, 55.86; %H, 5.02; %N, 14.99.

Example 15

2-Pentylthiazolo[4,5-*c*]quinoline-5N-oxide



Part A

Using the general method of Example 7 Part A, 2-methylthiazolo[4,5-*c*]quinoline (2.0 g, 10 mmol) was reacted first with lithium diisopropylamide (5.5 mL of 2M in benzene) and then with 1-iodobutane (1.8 mL) to provide 1.1 g of 2-pentylthiazolo[4,5-*c*]quinoline as a yellow solid, m.p. 62-64°C.


Part B

Peracetic acid (1.50 mL of 32% in acetic acid) was added to a suspension of 2-pentylthiazolo[4,5-*c*]quinoline (1.25 g) in methyl acetate (50 mL). The reaction mixture was heated at reflux for 6 hours. The reaction mixture was allowed to cool to ambient temperature and then it was diluted with dichloromethane and washed first with sodium bicarbonate and then with water. The organic layer was dried over magnesium sulfate and

then concentrated under vacuum to provide 1.20 g of a pale yellow solid. This material was recrystallized from ethyl acetate to provide 0.90 g of 2-pentylthiazolo[4,5-*c*]quinoline-5N-oxide as a white crystalline solid, m.p. 142-144°C. Analysis: Calculated for C₁₅H₁₆N₂OS: %C, 66.14; %H, 5.92; %N, 10.19; Found: %C, 65.63; %H, 5.83; %N, 10.28.

5

Example 16


2-Pentylthiazolo[4,5-*c*]quinolin-4-amine

Trichloroacetyl isocyanate (0.51 mL) was added to a solution of 2-pentylthiazolo[4,5-*c*]quinoline-5N-oxide (0.78 g) in dichloromethane (50 mL). The reaction mixture was stirred at ambient temperature for about 75 minutes and then concentrated under vacuum to provide crude N-(2-pentylthiazolo[4,5-*c*]quinolin-4-yl)trichloroacetamide. The amide was combined with a solution of ammonia in methanol (40 mL of 2M). Dichloromethane was added to bring all of the material into solution.

When the reaction was completed as indicated by thin layer chromatography, the reaction mixture was concentrated under vacuum. The residue was mixed with dichloromethane and sodium bicarbonate. The organic layer was separated, washed with sodium bicarbonate then with water, dried over magnesium sulfate and then concentrated under vacuum to provide a white solid. This material was recrystallized from ethyl acetate to provide 2-pentylthiazolo[4,5-*c*]quinolin-4-amine as an off-white crystalline solid, m.p. 119-121°C. Analysis: Calculated for C₁₅H₁₇N₃S: %C, 66.39; %H, 6.31; %N, 15.48; Found: %C, 66.21; %H, 6.35; %N, 15.39.

20

Example 17


2-Butylthiazolo[4,5-*c*]quinoline-5N-oxidePart A

- 5 Using the general method of Example 7 Part A, 2-methylthiazolo[4,5-*c*]quinoline (2.50 g, 12.5 mmol) was reacted first with lithium diisopropylamide (7.0 mL of 2M in benzene) and then with 1-iodopropane (3.0 g) to provide 1.19 g of 2-butylthiazolo[4,5-*c*]quinoline as a yellow oil.

Part B

- 10 Using the general method of Example 15 Part B, 2-butylthiazolo[4,5-*c*]quinoline (1.33 g) was oxidized with peracetic acid to provide 0.5 g of 2-butylthiazolo[4,5-*c*]quinoline-5N-oxide as a solid, m.p. 133-135°C.

Example 18


2-Butylthiazolo[4,5-*c*]quinolin-4-amine

15

Using the general method of Example 16, 2-butylthiazolo[4,5-*c*]quinoline-5N-oxide (0.50 g) was converted to the amide and then hydrolyzed to provide 0.25 g of 2-butylthiazolo[4,5-*c*]quinolin-4-amine as a yellow crystalline solid, m.p. 149-151°C.

- 20 Analysis: Calculated for C₁₄H₁₅N₃S: %C, 65.34; %H, 5.87; %N, 16.33; Found: %C, 64.88; %H, 5.84; %N, 16.03

Example 19

2-(1-Methylethyl)thiazolo[4,5-*c*]quinoline-5N-oxidePart A


- 5 Using the general method of Example 7 Part A, 2-methylthiazolo[4,5-*c*]quinoline (1.50 g, 7.5 mmol) was reacted first with lithium diisopropylamide (15.0 mL of 2M in benzene) and then with methyl iodide (2.4 mL) to provide 0.97 g of 2-(1-methylethyl)thiazolo[4,5-*c*]quinoline as a yellow oil.

Part B

- 10 Using the general method of Example 15 Part B, 2-(1-methylethyl)thiazolo[4,5-*c*]quinoline (0.95 g) was oxidized with peracetic acid to provide 0.84 g of 2-(1-methylethyl)thiazolo[4,5-*c*]quinoline-5N-oxide as a yellow solid, m.p. 161-162°C.

Example 20


- 15 2-(1-Methylethyl)thiazolo[4,5-*c*]quinolin-4-amine



- 20 Using the general method of Example 16, 2-(1-methylethyl)thiazolo[4,5-*c*]quinoline-5N-oxide (0.84 g) was converted to the amide and then hydrolyzed to provide 0.16 g of 2-(1-methylethyl)thiazolo[4,5-*c*]quinolin-4-amine as yellow needles, m.p. 163-165°C. Analysis: Calculated for C₁₃H₁₃N₃S: %C, 64.17; %H, 5.38; %N, 17.27; Found: %C, 63.49; %H, 5.36; %N, 17.09

Example 21

2-(2-Phenyl-1-ethenyl)thiazolo[4,5-c]quinoline-5N-oxide

Part A

- 5 Using the general method of Example 7 Part A, 2-methylthiazolo[4,5-c]quinoline (5.0 g, 25 mmol) was reacted first with lithium diisopropylamide (15.0 mL of 2M in benzene) and then with benzaldehyde (3.8 mL) to provide 5.3 g 1-phenyl-2-thiazolo[4,5-c]quinolin-2-yl-1-ethanol as a solid, m.p. 147-148°C.

Part B

- 10 Concentrated hydrochloric acid was added dropwise to a suspension of 1-phenyl-2-thiazolo[4,5-c]quinolin-2-yl-1-ethanol (2.16 g) in water (40 mL) until all of the solid had dissolved. The reaction mixture was heated on a steam bath during the addition; heating was continued until analysis by thin layer chromatography indicated that all of the starting material had reacted. The reaction mixture was allowed to cool to ambient temperature and a precipitate formed. The reaction mixture was neutralized with sodium carbonate.
- 15 Dichloromethane was added with stirring until all of the precipitate was in solution. The layers were separated and the aqueous layer was extracted with dichloromethane. The organic layers were combined, washed with water, dried over magnesium sulfate and then concentrated under vacuum to provide 2.2 g of a green solid. This material was
- 20 recrystallized from ethyl acetate to provide 1.55 g of 2-(2-phenyl-1-ethenyl)thiazolo[4,5-c]quinoline as a green crystalline solid. Analysis: Calculated for: C₁₈H₁₂N₂S: %C, 74.97; %H, 4.19; %N, 9.71; Found: %C, 74.89; %H, 4.17; %N, 9.72.

Part C


- 25 Peracetic acid (1.32 mL of 32% in acetic acid) was added to a suspension of 2-(2-phenyl-1-ethenyl)thiazolo[4,5-c]quinoline (1.20 g) in methyl acetate (50 mL). A precipitate formed. Ethanol was added to the reaction mixture until all of the precipitate was dissolved. The reaction mixture was heated at reflux overnight and then cooled to

ambient temperature. The resulting precipitate was isolated by filtration, dried and then recrystallized from methanol/dichloromethane to provide

2-(2-phenyl-1-ethenyl)thiazolo[4,5-*c*]quinoline-5N-oxide as a yellow solid, m.p. 268-270°C. Analysis: Calculated for: C₁₈H₁₂N₂OS: %C, 71.03; %H, 3.97; %N, 9.20; Found: %C, 69.94; %H, 3.87; %N, 9.05.

Example 22


2-(2-Phenyl-1-ethenyl)thiazolo[4,5-*c*]quinolin-4-amine



Using the general method of Example 16, 2-(2-phenyl-1-ethenyl)thiazolo[4,5-*c*]quinoline-5N-oxide (0.67 g) was converted to the trichloroacetamide then hydrolyzed to provide 0.43 g of 2-(2-phenyl-1-ethenyl)thiazolo[4,5-*c*]quinolin-4-amine as a yellow crystalline solid, m.p. 239-241°C. Analysis: Calculated for C₁₈H₁₃N₃S: %C, 71.26; %H, 4.32; %N, 13.85; Found: %C, 70.73; %H, 4.15; %N, 13.68.

Example 23

2-(2-Phenyl-1-ethyl)thiazolo[4,5-*c*]quinoline-5N-oxide



Part A

A small amount of catalyst (5% palladium on activated carbon) was added to a suspension of 2-(2-phenyl-1-ethenyl)thiazolo[4,5-*c*]quinoline (1.16 g, Example 21 Part B) in acetic acid (200 mL). The mixture was reduced on a Parr apparatus under a 50 psi (3.5 Kg/cm²) hydrogen atmosphere for 1 day. The reaction mixture was filtered to remove the catalyst. The filtrate was concentrated under vacuum. The residue was dissolved in

dichloromethane, washed with sodium bicarbonate then with water, dried over magnesium sulfate and then concentrated under vacuum to provide 0.88 g of 2-(2-phenyl-1-ethyl)thiazolo[4,5-c]quinoline as an oily solid.


Part B

- 5 Using the general method of Example 15 Part B, 2-phenylethylthiazolo[4,5-c]quinoline (0.90 g) was oxidized with peracetic acid to provide 0.63 g of 2-(2-phenyl-1-ethyl)thiazolo[4,5-c]quinoline-5N-oxide as an orange crystalline solid, m.p. 165-169°C. Analysis: Calculated for $C_{18}H_{14}N_2OS$: %C, 70.56; %H, 4.60; %N, 9.14; Found: %C, 69.59; %H, 4.50; %N, 9.04.

10

Example 24

2-(2-Phenyl-1-ethyl)thiazolo[4,5-c]quinolin-4-amine




- 15 Using the general method of Example 16, 2-(2-phenyl-1-ethyl)thiazolo[4,5-c]quinoline-5N-oxide (0.63 g) was converted to the trichloroacetamide then hydrolyzed to provide 0.21 g of 2-(2-phenyl-1-ethyl)thiazolo[4,5-c]quinolin-4-amine as a yellow crystalline solid, m.p. 158-159°C. Analysis: Calculated for $C_{18}H_{15}N_3S$: %C, 70.79; %H, 4.95; %N, 13.75; Found: %C, 70.29; %H, 4.90; %N, 13.66.

20

Example 25

2-Methyl-1-(thiazolo[4,5-c]quinolin-2-yl)-2-propanol-5N-oxide



Part A


Under an argon atmosphere, anhydrous tetrahydrofuran (150 mL) was added to a dried flask containing 2-methylthiazolo[4,5-*c*]quinoline (8.40 g). The reaction mixture was cooled to -78°C with a dry ice bath. Lithium diisopropylamide (23 mL of 2.0 M in benzene) was added dropwise. After about 50 minutes, acetone (5 mL) was added and the reaction mixture was allowed to warm to 0°C. After several hours the reaction was quenched with water, diluted with chloroform and then washed with water. The organic layer was dried over magnesium sulfate and then concentrated under vacuum. The residue was suspended in water (200 mL) and the mixture was heated. Hydrochloric acid (6N) was slowly added until all of the solid dissolved. Charcoal was added and the mixture was heated with stirring for about 5 minutes. The mixture was filtered to remove the charcoal. The filtrate was neutralized with sodium carbonate and then extracted with chloroform. The chloroform extract was washed several times with water, dried over magnesium sulfate and then concentrated under vacuum to provide 8.0 g of a light brown solid. This material was recrystallized from dichloromethane/hexanes to provide 5.0 g of 2-methyl-1-(thiazolo[4,5-*c*]quinolin-2-yl)-2-propanol as a yellow crystalline solid, m.p. 155-157°C. Analysis: Calculated for C₁₄H₁₄N₂OS: %C, 65.08; %H, 5.46; %N, 10.84; Found: %C, 64.97; %H, 5.33; %N, 10.90.

Part B

Peracetic acid (4.8 mL of 32% in acetic acid) was added to a suspension of 2-methyl-1-(thiazolo[4,5-*c*]quinolin-2-yl)-2-propanol (3.0 g) in methyl acetate (200 mL). The reaction mixture was heated at reflux overnight, cooled to ambient temperature and then concentrated under vacuum. The residue was dissolved in dichloromethane and then combined with sodium bicarbonate and stirred vigorously. The resulting precipitate was isolated by filtration and then dissolved in methanol/dichloromethane. This solution was concentrated under vacuum. The residue was combined with dichloromethane and then filtered to remove undissolved material. The filtrate was concentrated under vacuum to provide 2.6 g of the desired N-oxide. A small portion (0.2 g) was recrystallized from methanol/water to provide 2-methyl-1-(thiazolo[4,5-*c*]quinolin-2-yl)-2-propanol-5N-oxide as a solid, m.p. 187-189°C. Analysis: Calculated for C₁₄H₁₄N₂O₂S · 1/3 H₂O: %C, 59.98; %H, 5.27; %N, 9.99; Found: %C, 60.09; %H, 5.03; %N, 10.00.

Example 26

2-(4-Aminothiazolo[4,5-c]quinolin-2-yl)-1,1-dimethylethyl Carbamate




5 Trichloroacetyl isocyanate (3.2 mL) was added to a solution of 2-methyl-1-(thiazolo[4,5-c]quinolin-2-yl)-2-propanol-5N-oxide (2.4 g) in dichloromethane (250 mL). The reaction mixture was stirred at ambient temperature for 1 hour and then concentrated under vacuum. The residue was stirred with a solution of ammonia in methanol (150 mL of 2M) for 2 hours. The methanol was removed under vacuum. The residue was

10 suspended in a mixture of dichloromethane and ethyl acetate and then washed with sodium bicarbonate. The undissolved material was isolated by filtration, washed with water, washed with dichloromethane and then recrystallized from methanol/dichloromethane to provide 1.6 g of 2-(4-aminothiazolo[4,5-c]quinolin-2-yl)-1,1-dimethylethyl carbamate as a solid, m.p. 222-223°C. Analysis: Calculated for C₁₅H₁₆N₄O₂S: %C, 56.94; %H, 5.09;

15 %N, 17.70; Found: %C, 56.71; %H, 5.08; %N, 17.52.

Example 27

2-(Ethoxymethyl)thiazolo[4,5-c]quinoline-5N-oxide

20 Part A

Ethoxyacetyl chloride (6 mL, 53.8 mmol) was added to a suspension of 3-aminoquinoline-4-thiol (4.6 g, 26.1 mmol) in ethoxyacetic acid (50 mL). The reaction mixture was heated at 60°C overnight. The reaction mixture was concentrated under vacuum to remove a portion of the ethoxyacetic acid. The residue was combined with

water (100 mL) and a precipitate formed. The mixture was made basic with 50% sodium hydroxide. The precipitate was isolated by filtration, washed with water and then dried to provide 2-(ethoxymethyl)thiazolo[4,5-*c*]quinoline as a fluffy green solid.


Part B

5 Peracetic acid (1.0 mL of 32% in acetic acid) was added to a solution of 2-(ethoxymethyl)thiazolo[4,5-*c*]quinoline (1.0 g) in ethanol. The reaction mixture was stirred at ambient temperature for 1 week. The reaction mixture was concentrated under vacuum and then azeotroped with heptane to remove acetic acid. The residue was dissolved in dichloromethane, washed with sodium bicarbonate, washed with water, dried
10 over magnesium sulfate and then concentrated under vacuum. The residue was recrystallized from isopropanol to provide 2-(ethoxymethyl)thiazolo[4,5-*c*]quinoline-5N-oxide as a yellow crystalline solid, m.p. 138-140°C. Analysis: Calculated for $C_{13}H_{12}N_2O_2S$: %C, 59.98; %H, 4.65; %N, 10.76; Found: %C, 59.85; %H, 4.66; %N, 10.71.

15

Example 28

2-(Ethoxymethyl)thiazolo[4,5-*c*]quinoline-4-amine




Trichloroacetyl isocyanate (0.7 mL) was added to a solution of 2-(ethoxymethyl)thiazolo[4,5-*c*]quinoline-5N-oxide (1.0 g) in dichloromethane (50 mL).
20 The reaction mixture was stirred at ambient temperature for 2 hours and then concentrated under vacuum to provide N-(2-(ethoxymethyl)thiazolo[4,5-*c*]quinolin-4-yl)trichloroacetamide. The amide was taken up in methanol and then combined with 1 equivalent of sodium methoxide. The reaction mixture was stirred at ambient temperature
25 for 30 minutes and then concentrated under vacuum. The reaction was run a second time using 2 g of the N-oxide. The products were combined and recrystallized from isopropanol to provide 2.25 g of 2-(ethoxymethyl)thiazolo[4,5-*c*]quinoline-4-amine as

light yellow needles, m.p. 149-151°C. Analysis: Calculated for C₁₃H₁₃N₃OS: %C, 60.21; %H, 5.05; %N, 16.20; Found: %C, 59.86; %H, 4.97; %N, 16.16.

Example 29

5 2-(Methoxymethyl)thiazolo[4,5-*c*]quinoline-5N-oxide




Part A

Methoxyacetyl chloride (1.8 mL) was added to a mixture of 3-aminoquinoline-4-thiol (2.8 g) in methoxyacetic acid (15 mL). The reaction was heated at about 140°C for 1
10 hour and then allowed to cool to ambient temperature. The reaction mixture was diluted with a small amount of water, made basic with 10% sodium hydroxide and then extracted with dichloromethane (300 mL). The extract was washed with sodium bicarbonate, washed with water, dried over magnesium sulfate and then concentrated under vacuum to provide crude product as a dark oil. The oil was dissolved in dichloromethane and then
15 placed on a layer of silica gel. The silica gel was eluted with 1:1 hexane:ethyl acetate. The eluant was concentrated under vacuum to provide 2.3 g of 2-(methoxymethyl)thiazolo[4,5-*c*]quinoline as an orange solid.

Part B


Using the general method of Example 27 Part B, 2-(methoxymethyl)thiazolo[4,5-
20 *c*]quinoline (1.7 g) was oxidized to provide 1.8 g of 2-(methoxymethyl)thiazolo[4,5-*c*]quinoline-5N-oxide as yellow needles, m.p. 151-153°C. Analysis: Calculated for C₁₂H₁₀N₂OS: %C, 58.52; %H, 4.09; %N, 11.37; Found: %C, 57.95; %H, 3.98; %N, 11.3.

Example 30

2-(Methoxymethyl)thiazolo[4,5-*c*]quinolin-4-amine

Using the general method of Example 28, 2-(methoxymethyl)thiazolo[4,5-*c*]quinoline-5N-oxide (1.3 g) was reacted to form the trichloroacetamide and then hydrolyzed to provide 2-(methoxymethyl)thiazolo[4,5-*c*]quinolin-4-amine as light yellow needles, m.p. 183-185°C. Analysis: Calculated for C₁₂H₁₁N₃OS: %C, 58.76; %H, 4.52; %N, 17.13; Found: %C, 58.69; %H, 4.34; %N, 17.14.

Example 31


2-(2-Methylpropyl)thiazolo[4,5-*c*]quinoline-5N-oxidePart A

3-Aminoquinoline-4-thiol (4.6 g) was added to polyphosphoric acid (80 g). Isovaleric acid (3.5 mL) was added and the reaction mixture was heated at 140°C for 2 hours. The reaction mixture was poured into a mixture of ice and water (300 mL). The mixture was filtered through a layer of Celite® filter aid to remove some insoluble material. The filtrate was made alkaline with 50% sodium hydroxide while cooling with ice and then extracted with chloroform. The extract was washed with water, dried over magnesium sulfate and then concentrated under vacuum to provide an oil. The oil was dissolved in dichloromethane and then placed on a layer of silica gel and eluted with 1:1 ethyl acetate:hexanes. The eluant was concentrated under vacuum to provide 2-(2-methylpropyl)thiazolo[4,5-*c*]quinoline.

Part B


Using the general method of Example 27 Part B, 2-(2-methylpropyl)thiazolo[4,5-*c*]quinoline (5.2 g) was oxidized to provide 2.5 g of 2-(2-methylpropyl)thiazolo[4,5-*c*]quinoline-5N-oxide as a yellow solid.

5

Example 32**2-(2-Methylpropyl)thiazolo[4,5-*c*]quinolin-4-amine**

Using the general method of Example 28, 2-(2-methylpropyl)thiazolo[4,5-*c*]quinoline-5N-oxide (2.5 g) was reacted to form the trichloroacetamide and then hydrolyzed to provide 2-(2-methylpropyl)thiazolo[4,5-*c*]quinolin-4-amine as light yellow platelets, m.p. 123-125°C. Analysis: Calculated for C₁₄H₁₅N₃S: %C, 65.34; %H, 5.87; %N, 16.33; Found: %C, 64.87; %H, 5.79; %N, 16.18.

15

Example 33**2-Benzylthiazolo[4,5-*c*]quinoline-5N-oxide****Part A**

Thionyl chloride (1.5 g) was added dropwise to a cooled solution of phenyl acetic acid (2 g) in dichloromethane (10 mL). This mixture was allowed to stir at ambient temperature for 1 hour to provide a solution containing phenylacetyl chloride. Triethyl amine (4.3 mL) was added to a suspension of 3-aminoquinolin-4-ol in dichloromethane (10 mL) and the resulting mixture was cooled in an ice bath. The phenylacetyl chloride

20

solution was added dropwise to the cooled mixture. The reaction mixture was allowed to stir at ambient temperature overnight. The resulting thick oily precipitate was diluted with water (10 mL) and then stirred rapidly for 1 hour. The reaction mixture was filtered. Thin layer chromatography showed that both the isolated solid and the filtrate contained the
5 desired product. The filtrate was diluted with dichloromethane and water. The dichloromethane layer was separated, dried over magnesium sulfate and then concentrated under vacuum. The residue was combined with the previously isolated solid and recrystallized from 80:20 isopropanol:water to provide 1.3 g of N-(4-hydroxyquinolin-3-yl)phenylacetamide as needles, m.p. 253-255°C. Analysis: Calculated for: C₁₇H₁₄N₂O₂:
10 %C, 73.37; %H, 5.07; %N, 10.07; Found: %C, 73.16; %H, 5.03; %N, 10.07.

Part B

Phosphorous pentasulfide (1.6 g) was added to a suspension of N-(4-hydroxyquinolin-3-yl)phenylacetamide (1.0 g) in pyridine. The reaction mixture was heated at reflux until the reaction was complete. The reaction mixture was concentrated
15 under vacuum and then azeotroped with water to remove most of the pyridine. The residue was combined with water, neutralized with sodium carbonate and then extracted with dichloromethane. The extract was washed with water, dried over magnesium sulfate and then concentrated under vacuum to provide 2-benzylthiazolo[4,5-c]quinoline as a solid.


20 Part C

Using the general method of Example 27 Part B, 2-benzylthiazolo[4,5-c]quinoline (3.3 g) was oxidized to provide 2.1 g of 2-benzylthiazolo[4,5-c]quinoline-5N-oxide as a yellow solid, m.p. 185-186°C. Analysis: Calculated for C₁₇H₁₂N₂OS: %C, 69.84; %H, 4.14; %N, 9.58; Found: %C, 69.51; %H, 4.06; %N, 9.55

25

Example 34


2-Benzylthiazolo[4,5-c]quinolin-4-amine Hydrochloride



Trichloroacetyl isocyanate (1.2 mL, 10.3 mmol) was added to a solution of 2-benzylthiazolo[4,5-*c*]quinoline-5N-oxide (2.0 g, 6.8 mmol) in dichloromethane (100 mL). The reaction mixture was stirred at ambient temperature for 2 hours and then concentrated to provide crude N-(2-benzylthiazolo[4,5-*c*]quinolin-4-yl)trichloroacetamide. The amide was dissolved in methanol. Sodium methoxide (1 equivalent) was added. The reaction mixture was heated on a steam bath for 30 minutes and then allowed to cool to ambient temperature. The resulting precipitate was isolated by filtration and then suspended in a mixture of methanol and isopropanol. Hydrochloric acid (1 equivalent) was added and all of the solid dissolved initially. A white solid crystallized out. This material was isolated by filtration, washed with isopropanol and then dried to provide 1.5 g of 2-benzylthiazolo[4,5-*c*]quinolin-4-amine hydrochloride, m.p. 152-155°C. Analysis: Calculated for: C₁₇H₁₃N₃S · HCl: %C, 62.28; %H, 4.30; %N, 12.82; Found: %C, 62.05; %H, 4.23; %N, 12.82.

Example 35

8-Methyl-2-propylthiazolo[4,5-*c*]quinoline-5N-oxide



Part A

Catalyst (0.10 g of 10% platinum on carbon) was added to a solution of 6-methyl-3-nitroquinolin-4-ol (1 g) in ethanol (25 mL) and ammonium hydroxide (0.5 mL). The mixture was reduced on a Parr apparatus at ambient temperature under a hydrogen atmosphere. The reaction mixture was filtered to remove the catalyst and then concentrated under vacuum. The residue was combined with water and heated. Hydrochloric acid was added dropwise until all of the solid had dissolved. Activated carbon was added to the solution. The mixture was filtered. Hydrochloric acid (2 mL of 12N) was added to the filtrate. This recrystallization was run three times to provide 0.50 g of 3-amino-6-methylquinolin-4-ol hydrochloride, m.p. >310°C. Analysis: Calculated for

$C_{10}H_{10}N_2O \cdot HCl$: %C, 57.02; %H, 5.26; %N, 13.30; Found: %C, 56.92; %H, 5.16; %N, 13.24.

Part B

Triethylamine (11.46 mL) was added to a suspension of 3-amino-6-methylquinolin-4-ol hydrochloride in dichloromethane (400 mL). Butyryl chloride (4.46 mL) was added. The reaction mixture was heated on a steam bath for 30 minutes. The solution was diluted with sodium bicarbonate and then filtered. The filtrate was washed with bicarbonate and then concentrated under vacuum. The residue was recrystallized from isopropanol to provide 3-butyramido-6-methylquinolin-4-ol hemihydrate as a solid, m.p. 274-277°C. Analysis: Calculated for $C_{14}H_{16}N_2O_2 \cdot \frac{1}{2} H_2O$: %C, 66.39; %H, 6.76; %N, 11.06; Found: %C, 66.56; %H, 6.46; %N, 11.03.


Part C

Phosphorous pentasulfide (12.9 g) was added to a mixture of 3-butyramido-6-methylquinolin-4-ol hemihydrate (7.12 g) in pyridine. The reaction mixture was heated at reflux for 90 minutes, combined with a mixture of ice and sodium carbonate and then extracted with dichloromethane. The extract was concentrated under vacuum. The residue was diluted with toluene and then concentrated under vacuum to provide a crude solid. This material was purified using column chromatography eluting with 20% dichloromethane in ethyl acetate to provide 8-methyl-2-propylthiazolo[4,5-*c*]quinoline as a yellow solid.

Part D

Using the general method of Example 7 Part B, 8-methyl-2-propylthiazolo[4,5-*c*]quinoline (4.0 g) was oxidized using 3-chloroperoxybenzoic acid to provide 4.19 g of crude product which was recrystallized from isopropanol to provide 2.0 g of 8-methyl-2-propylthiazolo[4,5-*c*]quinoline-5N-oxide as a solid, m.p. 143-145°C. Analysis: Calculated for $C_{14}H_{14}N_2OS$: %C, 65.09; %H, 5.46; %N, 10.84; Found: %C, 64.86; %H, 5.40; %N, 10.88.


Example 36

8-Methyl-2-propylthiazolo[4,5-*c*]quinolin-4-amine

Using the general method of Example 28, 8-methyl-2-propylthiazolo[4,5-
5 *c*]quinoline-5N-oxide was converted to N-(8-methyl-2-propylthiazolo[4,5-*c*]quinolin-4-yl)trichloroacetamide and then hydrolyzed to provide 1.32 g of 8-methyl-2-propylthiazolo[4,5-*c*]quinolin-4-amine as a crystalline solid, m.p. 147-149°C. Analysis: Calculated for C₁₄H₁₅N₃S: %C, 63.54; %H, 5.87; %N, 16.33; Found: %C, 64.97; %H, 5.76; %N, 16.25.

10

Example 37

(4-Aminothiazolo[4,5-*c*]quinolin-2-yl)-methanolPart A

15 Triethylamine (7.3 mL) was added to a suspension of 3-aminoquinolin-4-ol (5 g) in dichloromethane (50 mL). The mixture was cooled in an ice bath and then acetoxyacetyl chloride (3 mL) was added dropwise. The reaction mixture was stirred at ambient temperature for 2 hours. The resulting thick precipitate was diluted with water (10 mL), stirred rapidly for 20 minutes and then isolated by filtration. Thin layer chromatography
20 indicated that both the solid and the filtrate contained the desired product. The filtrate was concentrated under vacuum. The residue was mixed with water and then filtered. The

combined solids were recrystallized from 80:20 isopropanol:water to provide N-(4-hydroxyquinolin-3-yl)acetoxyacetamide, m.p. 224-225°C.

Part B

Using the general method of Example 33 Part B, N-(4-hydroxyquinolin-3-yl)acetoxyacetamide (5.3 g) was reacted with phosphorous pentasulfide to provide 2.9 g of thiazolo[4,5-c]quinolin-2-ylmethyl acetate as a solid.

Part C


Using the general method of Example 27 Part B, thiazolo[4,5-c]quinolin-2-ylmethyl acetate (2.8 g) was oxidized with peracetic acid to provide thiazolo[4,5-c]quinolin-2-ylmethyl acetate-5N-oxide as a tan crystalline solid.

Part D

Trichloroacetyl isocyanate (0.65 mL) was added to a solution of thiazolo[4,5-c]quinolin-2-ylmethyl acetate-5N-oxide (1.0 g) in dichloromethane (50 mL). The reaction was stirred at ambient temperature for 2 hours and then concentrated under vacuum. The residue was dissolved in methanol. Sodium methoxide (1 equivalent) was added and the reaction mixture was stirred at ambient temperature overnight. The resulting precipitate was isolated by filtration and dried to provide 0.68 g of (4-aminothiazolo [4,5-c]quinolin-2-yl)methanol as a white solid, m.p. 247-249°C. Analysis: Calculated for C₁₁H₉N₃OS: %C, 57.13; %H, 3.92; %N, 18.17; Found: %C, 56.85; %H, 3.96; %N, 17.83.

Example 38

2-Methyloxazolo[4,5-c]quinolin-4-amine



Part A

3-Aminoquinolin-4-ol (6 g) was refluxed with acetic anhydride (8 eq.) until analysis by thin layer chromatography indicated that the reaction was complete. The reaction mixture was cooled, diluted with ice and water, made basic with 10% sodium

hydroxide and then extracted with dichloromethane. The extract was washed with water and brine, dried over magnesium sulfate and then concentrated under vacuum. The residue was purified by column chromatography (silica gel eluting with methanol/ethyl acetate) to provide 5.1 g of 2-methyloxazolo[4,5-c]quinoline.

5 Part B

A mixture of 2-methyloxazolo[4,5-c]quinoline (5.0 g), peracetic acid (5 eq.) and ethanol was stirred at ambient temperature. After 2 hours more peracetic acid (2 eq) was added and stirring was continued for 3 additional hours. The reaction mixture was concentrated under vacuum. The residue was azeotroped with heptane to provide 4.2 g of 2-methyloxazolo[4,5-c]quinoline-5N-oxide.


10 Part C

Trichloroacetyl isocyanate (3.6 mL) was slowly added to a cooled mixture of 2-methyloxazolo[4,5-c]quinoline-5N-oxide (4.0 g) and dichloromethane. The reaction mixture was stirred for several hours and then concentrated under vacuum to provide crude N-(2-methyloxazolo[4,5-c]quinolin-4-yl)trichloroacetamide. This material was combined with a solution of ammonia in methanol (2M) and stirred for 1 hour. The reaction mixture was concentrated under vacuum, diluted with water and then extracted with dichloromethane. The extract was washed with water and brine, dried over magnesium sulfate and then concentrated under vacuum. The residue was purified by column chromatography (silica gel eluting with ethyl acetate/hexane) to provide 1.2 g of 2-methyloxazolo[4,5-c]quinolin-4-amine as a solid, m.p. 195-197°C. Analysis: Calculated for C₁₁H₉N₃O: %C, 66.32; %H, 4.55; %N, 21.09; Found: %C, 65.96; %H, 4.44; %N, 20.68.

25

Example 39

2-Ethyloxazolo[4,5-c]quinolin-4-amine



Part A

3-Aminoquinolin-4-ol hydrochloride (6 g) was refluxed with propanoic anhydride (8 eq.) until analysis by thin layer chromatography indicated that the reaction was complete. The reaction mixture was cooled, diluted with ice and water, made basic with 10% sodium hydroxide and then extracted with dichloromethane. The extract was washed with water and brine, dried over magnesium sulfate and then concentrated under vacuum. The residue was purified by column chromatography (silica gel eluting with methanol/ethyl acetate) to provide 4.0 g of 2-ethyloxazolo[4,5-*c*]quinoline.

Part B


2-Ethyloxazolo[4,5-*c*]quinoline (3.5 g), peracetic acid (4.5 mL of 32% in acetic acid) and methyl acetate (40 mL) were combined and heated at 50°C for several hours. The reaction mixture was concentrated under vacuum. The residue was slurried with hexane and then filtered to provide 2.5 g of 2-ethyloxazolo[4,5-*c*]quinoline-5N-oxide as a solid.

Part C

Trichloroacetyl isocyanate (2 mL) was slowly added to a cooled mixture of 2-ethyloxazolo[4,5-*c*]quinoline-5N-oxide (2.5 g) and dichloromethane. The reaction mixture was stirred for several hours and then concentrated under vacuum to provide crude N-(2-ethyloxazolo[4,5-*c*]quinolin-4-yl)trichloroacetamide. This material was combined with a solution of ammonia in methanol (2M) and stirred for 1 hour. The reaction mixture was concentrated under vacuum, diluted with water and then extracted with dichloromethane. The extract was washed with water and brine, dried over magnesium sulfate and then concentrated under vacuum. The residue was purified by column chromatography (silica gel eluting with ethyl acetate/hexane) to provide 2-ethyloxazolo[4,5-*c*]quinolin-4-amine as a solid, m.p. 175-178°C. Analysis: Calculated for C₁₂H₁₁N₃O: %C, 67.59; %H, 5.20; %N, 19.71; Found: %C, 67.19; %H, 4.86; %N, 20.43.

Example 40

2-Butyloxazolo[4,5-c]quinoline-5N-oxide


Part A

- 5 3-Aminoquinolin-4-ol hydrochloride (1.97 g, 10.0 mmol), triethylamine (1.01 g, 10.1 mmol) and valeric anhydride (9.3 g, 50.0 mmol) were combined and then heated at reflux for 18 hours. The reaction mixture was cooled to ambient temperature and then poured over ice. The mixture was adjusted to pH 12 with 10% sodium hydroxide. The mixture was stirred until all of the ice melted and then it was extracted with diethyl ether.
- 10 The ether extracts were combined, dried over magnesium sulfate and then concentrated under vacuum to provide a tan solid. This material was purified by flash chromatography eluting with 3:2 ethyl acetate:dichloromethane to provide 1.45 g of 2-butyloxazolo[4,5-c]quinoline.

Part B


- 15 Peracetic acid (1.6 g, 6.8 mmol of 32% in acetic acid) was added with stirring to a solution of 2-butyloxazolo[4,5-c]quinoline (1.4 g, 6.2 mmol) in ethanol (50 mL). The reaction mixture was stirred at ambient temperature for 3 days and then quenched with saturated potassium carbonate solution. The layers were separated. The organic layer was concentrated under vacuum to provide a tan solid. This material was slurried with diethyl
- 20 ether and then filtered to provide 0.6 g of 2-butyloxazolo[4,5-c]quinolin-5N-oxide, m.p. 120-121°C. Analysis: Calculated for C₁₄H₁₄N₂O₂: %C, 69.41; %H, 5.82; %N, 11.56; Found: %C, 69.22; %H, 5.76; %N, 11.59.

Example 41

2-Butyloxazolo[4,5-*c*]quinolin-4-amine

Under a nitrogen atmosphere, trichloroacetyl isocyanate (0.6 g, 3.40 mmol) was
5 added with stirring to a solution of 2-butyloxazolo[4,5-*c*]quinolin-5N-oxide (0.55 g, 2.27
mmol) in anhydrous dichloromethane (20 mL). The reaction mixture was maintained at
ambient temperature for 2 hours and then concentrated under vacuum to provide crude N-
(2-butyloxazolo[4,5-*c*]quinolin-4-yl)trichloroacetamide as an oil. The oil was taken up in
methanol (25 mL). Sodium methoxide (0.49 g of 25%, 2.27 mmol) was added to the
10 solution. The reaction mixture was heated at reflux for 2 hours and then concentrated
under vacuum. The residue was taken up in ethyl acetate and washed with water. The
ethyl acetate layer was concentrated under vacuum to provide an orange solid. This
material was purified twice using flash chromatography eluting with ethyl acetate the first
time and with 30% dichloromethane in ethyl acetate the second time to provide 0.15 g of
15 2-butyloxazolo[4,5-*c*]quinolin-4-amine, m.p. 96-98°C. Analysis: Calculated for
C₁₄H₁₅N₃O: %C, 69.69; %H, 6.27; %N, 17.41; Found: %C, 69.23; %H, 6.06; %N, 17.07.

Example 42

2-Propylthiazolo[4,5-*c*]quinolin-4,8-diamine


Part A

Potassium nitrate (0.46 g, 4.52 mmol) was added to a solution of 2-propylthiazolo[4,5-*c*]quinolin-4-amine (1 g, 4.11 mmol, Example 12) in sulfuric acid (10 mL). The reaction mixture was stirred at ambient temperature for 30 minutes, then poured
5 onto ice, neutralized (pH=7) with ammonium hydroxide (150 mL) and then extracted with dichloromethane. The extract was washed with sodium bicarbonate, dried over magnesium sulfate and then concentrated under vacuum to provide 1 g of a yellow solid. This material was recrystallized from isopropanol/water to provide 0.84 g of 8-nitro-2-propylthiazolo[4,5-*c*]quinoline as a yellow solid, m.p. 228-230°C. Analysis: Calculated
10 for C₁₃H₁₂N₄O₂S: %C, 54.15; %H, 4.20; %N, 19.43; Found: %C, 54.22; %H, 4.05; %N, 19.04.

Part B

Catalyst (0.13 g of palladium on carbon) was added to a solution of 8-nitro-2-propylthiazolo[4,5-*c*]quinoline (1.31 g) in ethanol. The mixture was reduced on a Parr
15 apparatus under a hydrogen atmosphere. The reaction mixture was filtered to remove catalyst and the filter cake was washed with additional ethanol. The filtrate was concentrated under vacuum at 50°C and then oven dried under nitrogen to provide 2-propylthiazolo[4,5-*c*]quinolin-4,8-diamine as a yellow crystalline solid, m.p. 190-192°C, Analysis: Calculated for C₁₃H₁₄N₄S: %C, 60.44; %H, 5.46; %N, 21.69; Found: %C, 60.11;
20 %H, 5.45; %N, 21.96.

Example 43

2-Propyloxazolo[4,5-*c*]quinolin-4-amine25 Part A

A mixture of 3-aminoquinolin-4-ol hydrochloride (1.97 g, 10.0 mmol), butyric anhydride (3.15 g, 20 mmol) and pyridine (25 mL) was heated at reflux overnight. The


reaction mixture was cooled and then poured over ice. The mixture was made basic (pH 11) with 1 N sodium hydroxide and then it was extracted with diethyl ether (3 X 100 mL). A precipitate was removed by filtration. The ether extracts were combined, dried over magnesium sulfate and then concentrated to provide 1.1 g of 2-propyloxazolo[4,5-c]quinoline as an off white solid.

Part B


3-Chloroperoxybenzoic acid (1.0 eq. of 60%) was added with stirring to a solution of 2-propyloxazolo[4,5-c]quinoline (1.0 g, 4.7 mmol) in chloroform (30 mL). The reaction mixture was stirred at ambient temperature overnight and then quenched with a saturated potassium carbonate solution. The layers were separated. The aqueous layer was extracted with dichloromethane. The organic layers were combined and concentrated. The resulting crude product was purified by flash chromatography eluting with 8:2 ethyl acetate:dichloromethane to provide 1.0 g of 2-propyloxazolo[4,5-c]quinoline-5N-oxide as a tan solid.

Part C

Trichloroacetyl isocyanate (0.9 g, 5.25 mmol) was added with stirring to a solution of 2-propyloxazolo[4,5-c]quinoline-5N-oxide (0.8 g, 3.5 mmol) in dichloromethane (30 mL). The reaction mixture was stirred at ambient temperature for 2.5 hours and then concentrated under vacuum to provide crude N-(2-propyloxazolo[4,5-c]quinolin-4-yl)trichloroacetamide. The amide was dissolved in methanol (50 mL) and then combined with sodium methoxide (1.0 eq of 25% in methanol) and heated at reflux for 2 hours. The reaction mixture was concentrated under vacuum. The residue was taken up in diethyl ether and water. The ether layer was separated and concentrated to provide a tan solid. This material was purified by flash chromatography using two columns (The first was eluted with 8:2 ethyl acetate:dichloromethane; the second with 1:1 ethyl acetate:dichloromethane.) to provide 0.1 g of 2-propyloxazolo[4,5-c]quinolin-4-amine as a yellow powder, m.p. 159.0-160.0°C. Analysis: Calculated for C₁₃H₁₃N₃O: %C, 68.71; %H, 5.77; %N, 18.49; Found: %C, 68.03; %H, 5.77; %N, 18.14.

Example 448-Bromo-2-propylthiazolo[4,5-*c*]quinolin-4-amine

2-Propylthiazolo[4,5-*c*]quinolin-4-amine (1.0 g, 0.41 mmol) was combined with
5 acetic acid (15 mL) and heated to 60°C. Bromine (0.10 mL, 1.94 mmol) was added
dropwise and the reaction mixture was stirred at 60°C for 18 hours. The reaction mixture
was diluted with water and the resulting precipitate was isolated by filtration to provide
0.25 g of 8-bromo-2-propylthiazolo[4,5-*c*]quinolin-4-amine as a yellow solid, m.p. 177-
180°C. Analysis: Calculated for C₁₃H₁₂BrN₃S: %C, 48.46; %H, 3.75; %N, 13.04; Found:
10 %C, 47.98; %H, 3.95; %N, 12.70.

Example 457-Methyl-2-propylthiazolo[4,5-*c*]quinolin-4-amine

15

Part A

Diethyl ethoxymethylmalonate (37.8 mL, 187 mmol) and *m*-toluidine (20.0 mL,
187 mmol) were combined and heated at 100°C for about 3 hours. The reaction mixture
was allowed to cool to ambient temperature and it solidified. Dowtherm A (350 mL) was
20 added and the reaction mixture was heated at reflux for about 30 minutes. The reaction
mixture was allowed to cool to ambient temperature. The resulting precipitate was

isolated by filtration, rinsed with acetone and dried to provide 33 g of ethyl 4-hydroxy-7-methyl-3-quinolinecarboxylate as a tan powder.

Part B

Ethyl 4-hydroxy-7-methyl-3-quinolinecarboxylate (32 g, 138 mmol) was
5 suspended in sodium hydroxide (500 mL of 10% aqueous) and then heated at reflux for
about 30 minutes. The reaction mixture was allowed to cool to ambient temperature and
then it was acidified with concentrated hydrochloric acid. The resulting precipitate was
isolated by filtration, rinsed well with water and then oven dried to provide 4-hydroxy-7-
methyl-3-quinolinecarboxylic acid (28 g). A portion (2 g) was recrystallized twice from
10 N,N-dimethylformamide to give a fluffy white solid, m.p. 264-265°C. Analysis:
Calculated for $C_{11}H_9NO_3$: %C, 65.02; %H, 4.46; %N, 6.89; Found: %C, 65.22; %H, 4.42;
%N, 6.88.

Part C

4-Hydroxy-7-methyl-3-quinolinecarboxylic acid (32 g) was placed in a round
15 bottom flask and then heated in a Wood's metal bath at 310°C for several minutes until all
of the solids had melted into a light brown viscous liquid and bubbling had nearly ceased.
The reaction mixture was allowed to cool to ambient temperature. The crude solid was
recrystallized from ethyl acetate/ethanol to give 9.8 g of 7-methyl-4-quinolinol. During
the recrystallization a portion of the solid did not dissolve, this material was isolated by
20 filtration and then recrystallized to provide 1.1 g of 7-methyl-4-quinolinol as yellow-tan
plates, m.p. 233-235°C. Analysis: Calculated for $C_{10}H_9NO$: %C, 75.45; %H, 5.70; %N,
8.80; Found: %C, 75.23; %H, 5.54; %N, 8.76.

Part D

Nitric acid (6 mL of 70%) was slowly added to a hot (125°C) solution of 7-
25 methyl-4-quinolinol (10.5 g) in propionic acid (125 mL). The reaction mixture was stirred
for about 1.5 hours and then it was allowed to cool to ambient temperature. The resulting
precipitate was isolated by filtration, rinsed well with ethanol and water and then dried to
provide 6.9 g of 7-methyl-3-nitro-4-quinolinol as a pale yellow solid.

Part E

30 7-Methyl-3-nitro-4-quinolinol (11.8 g, 58 mmol), methanol (about 300 mL),
ammonium hydroxide (50 mL), and palladium on carbon (1 g of 10%) were combined.

The mixture was placed on a Parr apparatus under a 35-40 psi (2.4-2.8 Kg/cm²) hydrogen atmosphere for about 1 hour. The reaction mixture was filtered through a layer of Celite® filter agent and the filter cake was rinsed well with methanol. The filtrate was treated with charcoal and then concentrated under vacuum to provide a fluffy pale green solid. This material was triturated with acetonitrile to provide 8.5 g of 3-amino-7-methyl-4-quinolinol.

Part F

Under a nitrogen atmosphere, triethylamine (0.71 mL, 5.1 mmol) was added to a suspension of 3-amino-7-methyl-4-quinolinol (800 mg, 4.6 mmol) in dichloromethane (30 mL). Butyryl chloride (0.53 mL, 5.1 mmol) was added. The reaction mixture was stirred at ambient temperature for about 2 hours. Analysis by thin layer chromatography (silica gel eluting with 9:1 dichloromethane:methanol) showed starting material. The reaction mixture was heated to reflux and then inadvertently allowed to go dry over the course of about 30 minutes. More solvent was added and the reaction mixture was heated at reflux for an additional hour at which time thin layer chromatography showed no starting material. The resulting precipitate was isolated by filtration and then rinsed with dichloromethane and water to provide 650 mg of N-(4-hydroxy-7-methylquinolin-4-yl)butyramide as a pale pink-tan solid.

Part G

Under a nitrogen atmosphere, phosphorus pentasulfide (1.15 g, 2.6 mmol) was added to a mixture of N-(4-hydroxy-7-methylquinolin-4-yl)butyramide (630 mg, 2.6 mmol) in pyridine (20 mL). The reaction mixture was heated to reflux. The reaction mixture turned bright yellow and all of the solid went into solution. The reaction mixture was heated at reflux for about 2 hours and then allowed to cool to ambient temperature. The reaction mixture was extracted with water, aqueous sodium bicarbonate and dichloromethane. The organic layer was treated with saturated copper sulfate, dried with magnesium sulfate and then concentrated under vacuum to provide an oil. The oil was dried under high vacuum to provide 410 mg of 7-methyl-2-propylthiazolo[4,5-c]quinoline as an orange solid.

Part H

3-Chloroperoxybenzoic acid (2.4 g of 57-86%) was added to a mixture of 7-methyl-2-propylthiazolo[4,5-c]quinoline (2 g) and chloroform (100 mL). The reaction


mixture was allowed to stir at ambient temperature for 2 hours. Analysis by thin layer chromatography showed no starting material but did show two products. The reaction mixture was stirred at ambient temperature for an additional hour and then it was extracted with dichloromethane and aqueous sodium bicarbonate. The organic layer was dried with magnesium sulfate and then concentrated under vacuum to provide a yellow-orange oil. The oil was dried under high vacuum to provide 2.1 g of 7-methyl-2-propylthiazolo[4,5-c]quinolin-5N-oxide as a solid.

Part I

Under a nitrogen atmosphere, trichloroacetyl isocyanate (1.4 mL, 12.1 mmol) was added to a mixture of 7-methyl-2-propylthiazolo[4,5-c]quinolin-5N-oxide (2.1 g, 8.1 mmol) and dichloromethane (100 mL). The resulting dark brown solution was allowed to stir at ambient temperature for about 2 hours. The reaction mixture was concentrated under vacuum to provide N-(7-methyl-2-propylthiazolo[4,5-c]quinolin-4-yl)trichloroacetamide as an oil. The oil was combined with methanol and sodium methoxide (1.9 mL of 25% in methanol, 8.1 mmol) and then stirred at ambient temperature for 1 hour. The resulting precipitate was isolated by filtration and then recrystallized twice from isopropanol to provide 500 mg of 7-methyl-2-propylthiazolo[4,5-c]quinolin-4-amine as a yellow-tan powder, m.p. 186-187°C. Analysis: Calculated for $C_{14}H_{15}N_3S$: %C, 65.34; %H, 5.87; %N, 16.33; Found: %C, 64.95; %H, 5.77; %N, 16.08.

Example 46

2-Butyl-7-methyloxazolo[4,5-c]quinolin-4-amine



Part A

Under a nitrogen atmosphere, a mixture of 3-amino-7-methyl-4-quinolinol (5 g, 28.7 mmol) and valeric anhydride (28 mL, 143.5 mmol) was heated at reflux for about 20 hours. The reaction mixture was allowed to cool to ambient temperature, then it was made basic with 10% sodium hydroxide and stirred for an additional hour at ambient


temperature. The reaction mixture was extracted with dichloromethane. The extract was dried over magnesium sulfate and then concentrated under vacuum to provide a dark brown liquid. The liquid was purified by column chromatography (silica gel eluting with 3:2 ethyl acetate:dichloromethane) to provide 4.7 g of a dark brown, semi-solid. A portion
5 (about 700 mg) was purified by column chromatography (silica gel eluting with 95:5 dichloromethane:methanol) to provide 2-butyl-7-methyloxazolo[4,5-*c*]quinoline, m.p. 52-55°C. Analysis: Calculated for $C_{15}H_{16}N_2O$: %C, 74.97; %H, 6.71; %N, 11.66; Found: %C, 74.80; %H, 6.73; %N, 11.53.

Part B

10 3-Chloroperoxybenzoic acid (4.6 g of 57-86%) was added to a solution of 2-butyl-7-methyloxazolo[4,5-*c*]quinoline (3.9 g, 16.2 mmol) in chloroform (100 mL). The reaction mixture was allowed to stir at ambient temperature for 4 hours. The reaction mixture was washed with aqueous sodium bicarbonate, dried over magnesium sulfate and then concentrated under vacuum to provide 4.2 g of 2-butyl-7-methyloxazolo[4,5-
15 *c*]quinolin-5N-oxide as a dark brown-orange oil.

Part C

Under a nitrogen atmosphere, trichloroacetyl isocyanate (2.9 mL, 24 mmol) was added to a mixture of 2-butyl-7-methyloxazolo[4,5-*c*]quinolin-5N-oxide (4.2 g, 16 mmol) and dichloromethane (100 mL). The reaction mixture was allowed to stir at ambient
20 temperature for about 3 hours then it was concentrated under vacuum. The resulting residue was taken up in methanol and then combined with sodium methoxide (3.7 mL of 25 % in methanol, 16 mmol). The reaction mixture was stirred at ambient temperature overnight. The methanol was evaporated off and the resulting residue was purified by column chromatography (silica gel eluting with 95:5 dichloromethane:methanol) to
25 provide a brown solid. This solid was recrystallized from acetonitrile to provide 550 mg of 2-butyl-7-methyloxazolo[4,5-*c*]quinolin-4-amine as fine tan needles, m.p. 187-188°C. Analysis: Calculated for $C_{15}H_{17}N_3O + 0.1 H_2O$: %C, 70.07; %H, 6.74; %N, 16.34; Found: %C, 70.07; %H, 6.49; %N, 16.58.

Example 477-Methyl-2-propyloxazolo[4,5-*c*]quinolin-4-aminePart A

5 Under a nitrogen atmosphere a mixture of 3-amino-7-methyl-4-quinolinol (3.4 g, 20 mmol) and butyric anhydride (16 mL) were heated at reflux overnight. The reaction mixture was allowed to cool to ambient temperature and then it was poured over ice. The mixture was adjusted to pH 12 with 10% sodium hydroxide and then extracted with dichloromethane. The extract was dried over magnesium sulfate and then concentrated
10 under vacuum. The residue still contained anhydride so it was combined with 10 % sodium hydroxide and stirred for 1 hour at ambient temperature. The mixture was extracted with dichloromethane. The extract was dried over magnesium sulfate and then concentrated under vacuum to provide a brown oil. The oil was purified by column chromatography (silica gel eluting with 3:2 ethyl acetate:dichloromethane) to provide 3.1 g
15 of 7-methyl-2-propyloxazolo[4,5-*c*]quinoline as a light brown oil which solidified on standing, m.p. 65-68°C. Analysis: Calculated for C₁₄H₁₄N₂O: %C, 74.31; %H, 6.24; %N, 12.38; Found: %C, 73.69; %H, 6.07; %N, 12.15.

Part B

20 3-Chloroperoxybenzoic acid (3.8 g of 57-86%) was added to a solution of 7-methyl-2-propyloxazolo[4,5-*c*]quinoline (3 g) in chloroform (100 mL). The reaction mixture was allowed to stir at ambient temperature overnight. The reaction mixture was washed twice with sodium bicarbonate, dried over magnesium sulfate and then concentrated under vacuum to provide 3.1 g of 7-methyl-2-propyloxazolo[4,5-*c*]quinolin-5N-oxide as a pale orange solid.


Part C

25 Under a nitrogen atmosphere, trichloroacetyl isocyanate (2.3 mL, 19.2 mmol) was added to a solution of 7-methyl-2-propyloxazolo[4,5-*c*]quinolin-5N-oxide (3.1 g, 12.8

mmol) in dichloromethane (100 mL). The reaction mixture was allowed to stir at ambient temperature for 3 hours and then the solvent was removed under vacuum. Methanol (100 mL) was added to the resulting orange residue followed by sodium methoxide (2.9 mL of 25% in methanol, 12.8 mmol). The reaction mixture was stirred at ambient temperature overnight. The resulting precipitate was isolated by filtration and then recrystallized from isopropanol to provide 450 mg of 7-methyl-2-propyloxazolo[4,5-*c*]quinolin-4-amine as a white solid, m.p. 188-189°C. Analysis: Calculated for $C_{14}H_{15}N_3O + 0.2 H_2O$: %C, 68.66; %H, 6.34; %N, 17.16; Found: %C, 68.44; %H, 6.11; %N, 17.42.

10

Example 48
7-Fluoro-2-propyloxazolo[4,5-*c*]quinolin-4-amine



Part A

Under a nitrogen atmosphere, 3-fluoroaniline (50.0 g, 0.45 mol) and diethyl ethoxymethylmalonate (91 mL, 0.45 mol) were combined and heated at 100°C for 3 hours. The reaction was allowed to cool to ambient temperature and it solidified. Dowtherm A (200 mL) was added and the reaction mixture was heated at 240°C for 4 hours. The reaction mixture was allowed to cool to ambient temperature. The resulting precipitate was isolated by filtration, washed with hexane and then dried in a vacuum oven to provide 71.5 g of ethyl 7-fluoro-4-hydroxy-3-quinolinecarboxylate.

20

Part B

A suspension of ethyl 7-fluoro-4-hydroxy-3-quinolinecarboxylate (65 g, 0.28 mol) in 10% sodium hydroxide (250 mL) was heated at reflux for 3 hours during which time a solution was obtained. The reaction mixture was allowed to cool to ambient temperature and then it was filtered through filter paper under vacuum. The filtrate was acidified with concentrated hydrochloric acid. The resulting precipitate was collected, washed with

25

water and then dried to provide 53.5 g of 7-fluoro-4-hydroxy-3-quinolinecarboxylic acid as a white solid.

Part C

7-Fluoro-4-hydroxy-3-quinolinecarboxylic acid (25 g) was placed in a round bottom flask and heated to 330-350°C at which time carbon dioxide liberation commenced and the material began to liquefy. After about 2 minutes an additional 25 g of 7-fluoro-4-hydroxy-3-quinolinecarboxylic acid was added. The heating was continued for an additional 4 to 6 minutes at which time there was no further evolution of carbon dioxide. The solution was allowed to cool to ambient temperature. The resulting solid was isolated by filtration to provide 35.6 g of 7-fluoro-4-quinolinol as a pink solid.

Part D

Nitric acid (20 mL of 70%) was added to a hot (125°C) solution of 7-fluoro-4-quinolinol (35 g, 214 mmol) in propionic acid (200 mL). The reaction mixture was stirred at 125°C for about 1.5 hours and then allowed to cool to ambient temperature. The resulting yellow precipitate was isolated by filtration, rinsed with water and then with ethanol, and then recrystallized from N,N-dimethylformamide/water to provide 18 g of 7-fluoro-3-nitro-4-quinolinol.

Part E

A mixture containing 7-fluoro-3-nitro-4-quinolinol (17 g, 81.7 mmol), ammonium hydroxide (80 mL), methanol (200 mL) and palladium on carbon (1 g of 10% was placed on a Parr apparatus under a hydrogen atmosphere of about 30 psi (2.1 Kg/cm²) for 1 hour. The reaction mixture was filtered to remove the catalyst. The filtrate was treated with charcoal then concentrated under vacuum to provide a dark tan solid which turned a very dark brown upon oven drying. The solid was dissolved in methanol then hydrochloric acid in diethyl ether was added. A gray precipitate formed almost immediately. The suspension was stirred at ambient temperature for several hours. The precipitate was isolated by filtration and rinsed well with ether to provide 6.6 g of 3-amino-7-fluoro-4-quinolinol hydrochloride.

Part F

Under a nitrogen atmosphere, 3-amino-7-fluoro-4-quinolinol hydrochloride (3.4 g, 19.1 mmol), triethylamine (2.9 mL, 21.0 mmol) and butyric anhydride (15.6 mL, 95.5

mmol) were combined and heated at reflux for about 18 hours. The reaction mixture was poured over ice and made basic to about pH 12 with 10% sodium hydroxide. The resulting suspension was stirred until all of the ice had melted then it was extracted with dichloromethane. The extract was dried over magnesium sulfate and then concentrated under vacuum to provide an oil. The oil was purified by column chromatography (silica gel eluting initially with dichloromethane and then with 9:1 dichloromethane: methanol) to provide 2.6 g of 7-fluoro-2-propyloxazolo[4,5-*c*]quinoline as a light brown solid.

Part G

7-Fluoro-2-propyloxazolo[4,5-*c*]quinoline (2.6 g, 11.3 mmol), 3-chloroperoxybenzoic acid (3.3 g of 57-86%) and chloroform (90 mL) were combined and stirred at ambient temperature for about 3 hours. Analysis by thin layer chromatography (silica gel eluting with 95:5 dichloromethane:methanol) showed starting material. An additional 0.5 equivalent of 3-chloroperoxybenzoic acid was added and the reaction was stirred at ambient temperature for another 2 hours at which time thin layer chromatography showed no starting material. The reaction mixture was diluted with dichloromethane and then washed twice with sodium bicarbonate. The organic layer was dried over magnesium sulfate and then concentrated under vacuum to provide 2.8 g of 7-fluoro-2-propyloxazolo[4,5-*c*]quinolin-5N-oxide as a yellow-orange oily solid.


Part H

Under a nitrogen atmosphere, trichloroacetyl isocyanate (2.0 mL, 17.0 mmol) was added to a solution of 7-fluoro-2-propyloxazolo[4,5-*c*]quinolin-5N-oxide (2.8 g, 11.3 mmol) in dichloromethane (50 mL). The reaction mixture was stirred at ambient temperature for 3 hours and then the dichloromethane was removed under vacuum. The residue was dissolved in methanol and then combined with sodium methoxide (2.4 mL of 25 % in methanol, 11.3 mmol). The reaction mixture was stirred at ambient temperature overnight then filtered to remove a small amount of solid material. The filtrate was concentrated under vacuum. The residue was dissolved in dichloromethane, dried over magnesium sulfate and then concentrated under vacuum to provide a brown oil. The oil was purified by column chromatography (silica gel eluting with 95:5 dichloromethane:methanol) to provide a light brown sticky solid. This material was recrystallized from acetonitrile to provide 200 mg of 7-fluoro-2-propyloxazolo[4,5-

c]quinolin-4-amine as a rust colored powder, m.p. 184-187°C. Analysis: Calculated for $C_{13}H_{12}FN_3O$: %C, 63.67; %H, 4.93; %N, 17.13; Found: %C, 63.43; %H, 4.57; %N, 16.74.

Example 49

7-Fluoro-2-propylthiazolo[4,5-c]quinolin-4-amine

Part A

Under a nitrogen atmosphere, triethylamine (6.4 mL, 46.2 mmol) was added to a suspension of 3-amino-7-fluoro-4-quinolinol hydrochloride (3 g, 14.0 mmol) in tetrahydrofuran (50 mL). Butyryl chloride (1.6 mL, 15.4 mmol) was added dropwise at ambient temperature. The reaction mixture was stirred at ambient temperature for 4 hours. Aqueous sodium bicarbonate was added and the reaction mixture was allowed to stir at ambient temperature for about 1 hour. The resulting biphasic mixture was filtered to remove solids. The solids were rinsed with diethyl ether to provide a slightly pink powder. The tetrahydrofuran layer was concentrated under vacuum to provide a dark pink solid. This solid was triturated with ether and then oven dried. The solids were combined to provide 3.0 g of N-(7-fluoro-4-hydroxyquinolin-3-yl)butanamide. A 300 mg portion was recrystallized from ethyl acetate/ethanol to provide a light gray fluffy solid, m.p. 306-308°C. Analysis: Calculated for $C_{13}H_{13}FN_2O_2$: %C, 62.90; %H, 5.28; %N, 11.28; Found: %C, 62.95; %H, 5.34; %N, 11.14.

Part B

Under a nitrogen atmosphere, phosphorus pentasulfide (4.7 g, 10.5 mmol) was added to a mixture of N-(7-fluoro-4-hydroxyquinolin-3-yl)butanamide (2.6 g, 10.5 mmol) and pyridine (80 mL). The reaction mixture was heated at reflux for 2 hours and then allowed to cool to ambient temperature. The reaction mixture was extracted with sodium bicarbonate/dichloromethane. The organic layer was separated, washed twice with water, dried over magnesium sulfate and then concentrated under vacuum to provide a rust

colored solid. This material was recrystallized from methanol to provide 1.8 g of 7-fluoro-2-propylthiazolo[4,5-c]quinoline as rust colored platelike needles.


Part C

3-Chloroperoxybenzoic acid (2.1 g of 57-86%) was added to a solution of 7-fluoro-2-propylthiazolo[4,5-c]quinoline (1.8 g, 7.3 mmol) in chloroform (50 mL). The reaction mixture was stirred at ambient temperature. Analysis by thin layer chromatography (silica gel eluting with 95:5 dichloromethane:methanol) showed starting material so an additional 0.5 equivalent of 3-chloroperoxybenzoic acid was added. The reaction mixture was stirred at ambient temperature overnight. The reaction mixture was washed twice with sodium bicarbonate, dried over magnesium sulfate and then concentrated under vacuum to provide 1.8 g of 7-fluoro-2-propylthiazolo[4,5-c]quinolin-5N-oxide as a pale orange solid.

Part D

Under a nitrogen atmosphere, trichloroacetyl isocyanate (1.2 mL, 10.4 mmol) was added to a mixture of 7-fluoro-2-propylthiazolo[4,5-c]quinolin-5N-oxide (1.8 g, 6.9 mmol) and dichloromethane (50 mL). The reaction mixture was stirred at ambient temperature for 2 hours and then concentrated under vacuum to provide N-(7-fluoro-2-propylthiazolo[4,5-c]quinolin-4-yl)trichloroacetamide as an orange oil. The oil was dissolved in methanol and then combined with sodium methoxide (1.5 mL of 25 wt% in methanol). The reaction mixture was allowed to stir at ambient temperature for 3 hours. The resulting precipitate was isolated by filtration and then recrystallized first from acetonitrile and then from methanol to provide 1.1 g of 7-fluoro-2-propylthiazolo[4,5-c]quinolin-4-amine as a tan powder, m.p. 192.5-193.5°C. Analysis: Calculated for $C_{13}H_{12}FN_3S$: %C, 59.75; %H, 4.63; %N, 16.08; Found: %C, 59.55; %H, 4.69; %N, 16.12.

Example 50
2-Propyl-7-(trifluoromethyl)thiazolo[4,5-c]quinolin-4-amine



Part A

5 Under a nitrogen atmosphere, a mixture of 3-(trifluoromethyl)aniline (40 mL, 0.32 mmol) and diethyl ethoxymethylmalonate was heated at 100°C for 3 hours. The reaction mixture was allowed to cool to room temperature at which time the solution solidified to provide 102 g of diethyl 2-{{3-(trifluoromethyl)anilino}methylene}malonate as a cream colored solid.

10 Part B

Under a nitrogen atmosphere, a mixture of 2-{{3-(trifluoromethyl)anilino}methylene}malonate (80 g, 0.24 mol) and Dowtherm A was heated to 240°C and then stirred for 3 hours. The reaction mixture was allowed to cool to ambient temperature and then stirred for 16 hours. The solids were isolated by filtration then washed with hexane to provide 47.5 g of ethyl 4-hydroxy-7-(trifluoromethyl)-3-quinolinecarboxylate as an off-white solid.

Part C

A mixture of ethyl 4-hydroxy-7-(trifluoromethyl)-3-quinolinecarboxylate (43.4 g, 0.521 mol) and 10% sodium hydroxide (150 mL) was heated to reflux. Most of the ester did not dissolve so methanol (150 mL) was added over the course of an hour to facilitate dissolution. After refluxing for 2 hours a solution was obtained. The solution was refluxed for an additional 2 hours and then allowed to cool to ambient temperature overnight. The methanol was removed under reduced pressure and the resulting aqueous solution was acidified with concentrated hydrochloric acid. The resulting precipitate was isolated by filtration, washed with water and then dried in a vacuum oven at 120°C for 24

hours to provide 38.5 g of 4-hydroxy-7-(trifluoromethyl)-3-quinolinecarboxylic acid as a white solid.

Part D

5 A round bottom flask was charged with 4-hydroxy-7-(trifluoromethyl)-3-quinolinecarboxylic acid (34.1 g, 0.132 mol) and then heated in a Wood's metal bath for 5 minutes during which time carbon dioxide evolution was observed and the material changed from a solid to a liquid. After 5 minutes no further gas evolution was noted so the flask was removed from the bath and allowed to cool to ambient temperature. The resulting solid was isolated by filtration to provide 27.75 g of 7-(trifluoromethyl)-4-quinolinol.
10

Part E

A mixture of 7-(trifluoromethyl)-4-quinolinol (22.7 g, 0.106 mol) and propionic acid (106 mL) was heated to 120°C. Nitric acid (10 mL of 70%) was added dropwise and heating was continued for an additional 2 hours. The reaction mixture was allowed to cool
15 to ambient temperature. The resulting precipitate was isolated by filtration then washed with water and diethyl ether to provide 13.3 g of 3-nitro-7-(trifluoromethyl)-4-quinolinol as an off-white solid.

Part F

A Parr flask was charged with methanol (40 mL), ammonium hydroxide (10 mL),
20 3-nitro-7-(trifluoromethyl)-4-quinolinol (12.8 g, 49.6 mmol) and palladium on carbon (1.0 g of 10%). The mixture was placed on a Parr apparatus under a hydrogen atmosphere at 40 psi (2.8 Kg/cm²) for 4 hours. The mixture was filtered and the catalyst was washed with methanol and dichloromethane. The combined organics were concentrated under vacuum to provide a green solid. The solid was dissolved in methanol and then combined
25 with 1 N hydrochloric acid in anhydrous diethyl ether (150 mL). A precipitate formed almost immediately. The reaction mixture was allowed to stir for 16 hours. The precipitate was isolated by filtration, washed with diethyl ether and then dried in a vacuum oven at 80°C to provide 9.3 g of 3-amino-7-(trifluoromethyl)-4-quinolinol hydrochloride as an off-white solid.

Part G

Butyryl chloride (1.5 mL, 14.5 mmol) was added dropwise to a mixture of 3-amino-7-(trifluoromethyl)-4-quinolinol hydrochloride (3.5 g, 13.2 mmol), triethylamine (6.1 mL, 43.6 mmol) and anhydrous tetrahydrofuran (30 mL). The reaction mixture was allowed to stir for 16 hours. A small amount of aqueous sodium bicarbonate was added and the reaction mixture was stirred for 0.5 hours. The tetrahydrofuran was removed under vacuum. The resulting solid was stirred with diethyl ether, isolated by filtration, washed with water and diethyl ether, and then dried in a vacuum oven at 80°C overnight to provide 3.3 g of N-[4-hydroxy-7-(trifluoromethyl)quinolin-3-yl]butanamide as a cream colored solid.

Part H

A mixture of N-[4-hydroxy-7-(trifluoromethyl)quinolin-3-yl]butanamide (3.0 g, 10.05 mmol), phosphorous pentasulfide (4.5 g, 10.05 mmol) and pyridine (30 mL) was heated at reflux for 6 hours. The solution was allowed to cool to ambient temperature and then it was diluted with dichloromethane and aqueous sodium bicarbonate. The organic layer was separated, washed with water and with brine, dried over magnesium sulfate and then concentrated under vacuum to provide a yellow solid. This material was triturated with hexane and then isolated by filtration to provide 1.7 g 2-propyl-7-(trifluoromethyl)thiazolo[4,5-c]quinoline as a tan solid. The hexane filtrate was concentrated to give 0.6 g of additional product as a yellow solid.

Part I

3-Chloroperoxybenzoic acid (1.93 g, 6.88 mol) was added to a mixture of 2-propyl-7-(trifluoromethyl)thiazolo[4,5-c]quinoline (2.0 g, 6.75 mmol) in chloroform (30 mL). The resulting solution was allowed to stir for 24 hours. The reaction mixture was diluted with aqueous sodium bicarbonate and then extracted with dichloromethane. The organic layer was separated, dried over magnesium sulfate and then concentrated under vacuum to provide 1.98 g of 2-propyl-7-(trifluoromethyl)thiazolo[4,5-c]quinolin-5N-oxide as a yellow solid.


Part J

Under a nitrogen atmosphere, trichloroacetyl isocyanate (0.75 mL, 6.24 mmol) was added to a mixture of 2-propyl-7-(trifluoromethyl)thiazolo[4,5-c]quinolin-5N-oxide (1.3 g,

4.16 mmol) and anhydrous dichloromethane (20 mL). The resulting solution was allowed to stir at ambient temperature for 16 hours. The solvent was removed under reduced pressure. The resulting residue was dissolved in methanol (40 mL) then combined with sodium methoxide (1.43 mL of 25% in methanol, 6.24 mmol). The resulting solution was allowed to stir at ambient temperature for 16 hours by which time a precipitate had formed. The precipitate was isolated by filtration, washed with a small amount of methanol and then dried for 16 hours in a vacuum oven at 80°C to provide 0.96 g of 2-propyl-7-(trifluoromethyl)thiazolo[4,5-c]quinolin-4-amine as a white solid, m.p. 215-16°C. Analysis: Calculated for C₁₄H₁₂F₃N₃S: %C, 54.01; %H, 3.89; %N, 13.50; Found: %C, 53.82; %H, 3.66; %N, 13.37.

Example 51

2-(Methylsulfonyl)thiazolo[4,5-c]quinolin-5N-oxide



Part A

N⁴-(2-methylpropyl)quinoline-3,4-diamine (5.4 g, 25 mmol) was combined with carbon disulfide (9 mL, 150 mmol) and ethanol (55 mL) and then heated at reflux on a steam bath for 2 hours. The resulting precipitate was isolated by filtration, washed with ethanol and then air dried to provide 4.4 g of crude product. A portion (1 g) was dissolved in hot dilute sodium hydroxide and then reprecipitated with acetic acid. The precipitate was isolated by filtration while still hot, washed with hexane and then air dried to provide thiazolo[4,5-c]quinoline-2-thiol as a solid, m.p. 282-284°C. Analysis: Calculated for C₁₀H₆N₂S₂: %C, 55.02; %H, 2.77; %N, 12.83; Found: %C, 54.96; %H, 2.69; %N, 12.74.

Part B

Sodium methoxide (15.8 mL of 25% in methanol, 69 mmol) and methyl iodide (3.9 mL, 63 mmol) were added to a solution of thiazolo[4,5-c]quinoline-2-thiol (13.65 g, 63 mmol) in methanol (160 mL). The reaction mixture was heated on a steam bath for 1 hour. The solvent was removed under vacuum. The resulting light green-yellow solid was


slurried with water, isolated by filtration, and washed with water to provide 9.8 g of crude product. A portion (1 g) was recrystallized from methanol to provide 2-(methylthio)thiazolo[4,5-*c*]quinoline as a solid, m.p. 116-119°C. Analysis: Calculated for C₁₁H₈N₂S₂: %C, 56.87; %H, 3.47; %N, 12.06; Found: %C, 57.09; %H, 3.57; %N, 12.04.

5 Part C

Peracetic acid (27.8 mL of 32%, 132 mmol) was added to a mixture of 2-(methylthio)thiazolo[4,5-*c*]quinoline (7.7 g, 33 mmol) and acetic acid (100 mL). The reaction mixture was heated at about 60°C for about 4 hours and then at ambient temperature overnight. The resulting yellow precipitate was isolated by filtration to give 10 5.6 g of crude product. The filtrate was concentrated under vacuum then the residue was diluted with toluene (100 mL). The toluene was removed under vacuum to provide an additional 4 g of crude product. A portion (1 g) was recrystallized from N,N-dimethylformamide to provide 2-(methylsulfonyl)thiazolo[4,5-*c*]quinolin-5N-oxide as a yellow solid, m.p. 245-247°C. Analysis: Calculated for C₁₁H₈N₂O₃S₂: %C, 47.13; %H, 15 2.88; %N, 9.99; Found: %C, 47.08; %H, 3.08; %N, 10.14.

Example 52


2-(4-Morpholino)thiazolo[4,5-*c*]quinoline-5N-oxide



20 2-(Methylsulfonyl)thiazolo[4,5-*c*]quinolin-5N-oxide (2.5 g, 8.9 mmol) and morpholine (~50 mL) were combined and then heated on a steam bath for 9 hours. The resulting precipitate was isolated by filtration to provide 0.9 g of crude product as a yellow solid. The filtrate was cooled in an ice bath. The resulting precipitate was isolated by filtration to give 0.8 g of crude product as a yellow solid. The two crops were combined 25 and then a portion (0.5 g) was recrystallized from methanol to provide 2-(4-morpholino)thiazolo[4,5-*c*]quinoline-5N-oxide as a solid, m.p. 241-242°C. Calculated for C₁₄H₁₃N₃O₂S: %C, 58.52; %H, 4.56; %N, 14.62; Found: %C, 58.24; %H, 4.38; %N, 14.43.

Example 53

2-(4-Morpholino)thiazolo[4,5-c]quinolin-4-amine




5

Ammonium hydroxide (18 mL) was added to a mixture of 2-(4-morpholino)thiazolo[4,5-c]quinoline-5N-oxide (1.2 g, 4.2 mmol) and dichloromethane (24 mL). The mixture was cooled and then tosyl chloride (0.88 g, 4.6 mmol) in dichloromethane (10 mL) was slowly added. The reaction mixture was warmed to ambient temperature and then stirred overnight. The organic phase was separated, washed with aqueous sodium bicarbonate, dried over magnesium sulfate and then concentrated under vacuum to provide crude product as a yellow solid. This material was purified by column chromatography then dissolved in hydrochloric acid and reprecipitated with sodium hydroxide. The precipitate was isolated by filtration and then recrystallized twice from methanol to provide 0.26 g of 2-(4-morpholino)thiazolo[4,5-c]quinolin-4-amine as a solid, m.p. 225-227°C. Analysis: Calculated for C₁₄H₁₄N₄OS: %C, 58.72; %H, 4.93; %N, 19.57; Found: %C, 58.47; %H, 4.63; %N, 19.23.

15

Example 54

2-(1-Pyrrolidino)thiazolo[4,5-c]quinolin-4-amine




20

Part A

2-(Methylsulfonyl)thiazolo[4,5-*c*]quinolin-5N-oxide (2.5 g, 8.9 mmol) and pyrrolidine (~70 mL) were combined and then refluxed on a steam bath for 3 days. The resulting yellow precipitate was isolated by filtration to provide 0.4 g of 2-(1-pyrrolidino)thiazolo[4,5-*c*]quinoline-5N-oxide. The filtrate was cooled in an ice bath. The resulting precipitate was isolated by filtration to provide 0.7 g of 2-(1-pyrrolidino)thiazolo[4,5-*c*]quinoline-5N-oxide as a yellow solid. The two crops were combined.

Part B

Ammonium hydroxide (12 mL) was added to a mixture of 2-(1-pyrrolidino)thiazolo[4,5-*c*]quinoline-5N-oxide (0.8 g, 2.95 mmol) and dichloromethane (50 mL). The mixture was cooled and then tosyl chloride (0.6 g, 3.2 mmol) in dichloromethane (10 mL) was slowly added. The reaction mixture was warmed to ambient temperature and then stirred overnight. The organic phase was separated, washed with saturated aqueous sodium bicarbonate and then concentrated under vacuum to provide crude product as a yellow solid. This material was purified by flash column chromatography then slurried with hot methanol, cooled and isolated by filtration to provide 0.14 g of 2-(1-pyrrolidino)thiazolo[4,5-*c*]quinolin-4-amine as a solid, m.p. 259-261°C. Analysis: Calculated for C₁₄H₁₄N₄S: %C, 62.20; %H, 5.22; %N, 20.49; Found: %C, 61.76; %H, 5.25; %N, 20.72.

Example 552-Propylthiazolo[4,5-*c*]quinolin-4-amine Xinofate


2-Propylthiazolo[4,5-*c*]quinolin-4-amine (3.0 g, 12.3 mmol) and 1-hydroxy-2-naphthoic acid (2.3 g, 12.3 mmol) were separately dissolved in methanol with the use of dichloromethane if necessary. The two solutions were combined and the resulting solution

was reduced in volume. The resulting precipitate was isolated by filtration to provide 3.6 g of 2-propylthiazolo[4,5-*c*]quinolin-4-amine xinofoate as a colorless crystalline solid, m.p. 185-189°C (decomposed). Analysis: Calculated for $C_{24}H_{21}N_3O_3S$: %C, 66.80; %H, 4.91; %N, 9.74; Found: %C, 66.71; %H, 5.07; %N, 9.78.

5

Example 56

2-Propylthiazolo[4,5-*c*]quinolin-4-amine 3-Hydroxy-2-naphthoate




10 A solution of 3-hydroxy-2-naphthoic acid (1.9 g, 10 mmol) in methanol (30 mL) was added to a solution of 2-propylthiazolo[4,5-*c*]quinolin-4-amine (2.4 g, 10 mmol) in hot methanol (70 mL). A precipitate formed immediately. The mixture was heated an additional 5 minutes and then allowed to cool to ambient temperature. The precipitate was isolated by filtration, washed with methanol and dried to provide 4.0 g of product as a tan powder. This material was recrystallized from methanol/dichloromethane to provide 3.2 g of 2-propylthiazolo[4,5-*c*]quinolin-4-amine 3-hydroxy-2-naphthoate as a white powder. Calculated for $C_{24}H_{21}N_3O_3S$: %C, 66.80; %H, 4.91; %N, 9.74; Found: %C, 66.28; %H, 4.92; %N, 9.59.

20

Example 57

2-Butylthiazolo[4,5-*c*][1,5]naphthyridine-5N-oxide



Part A

25 A mixture containing 3-nitro[1,5]naphthyridin-4-ol (7.5 g), methanol (200 mL), ammonium hydroxide (50 mL) and 5% platinum on carbon (0.75 g) was placed on a Parr

apparatus for 6 hours. The reaction mixture was filtered to remove catalyst and then filtered a second time using Celite® filter aid. The filtrate was concentrated under vacuum to provide 6.1 g of 3-amino[1,5]naphthyridin-4-ol as a brown solid.

Part B

5 Valeryl chloride (4.3 g, 35 mmol) was added dropwise to a suspension of 3-amino[1,5]naphthyridin-4-ol (5.2 g, 32 mmol) in pyridine (100 mL). The reaction mixture was heated at reflux for 2 hours. The pyridine was removed. The resulting residue was taken up in hot water and then allowed to cool. The resulting gray precipitate was isolated by filtration, washed well with hot water and then oven dried to provide 2.3 g of N-(4-
10 hydroxy[1,5]naphthyridin-3-yl)pentamide as a gray solid.

Part C

Phosphorous pentasulfide (4.2 g, 9.4 mmol) was added to a suspension of N-(4-hydroxy[1,5]naphthyridin-3-yl)pentamide (2.3 g, 9.4 mmol) in pyridine (150 mL). The reaction mixture was heated at reflux for 2 hours. The pyridine was removed. The
15 resulting residue was taken up in a mixture of water, 10% sodium carbonate and 10% sodium hydroxide (an amount sufficient to adjust the pH to >8) and then extracted twice with dichloromethane. The dichloromethane extracts were combined, washed with brine, dried and then concentrated under vacuum. The residue was diluted with toluene and then concentrated under vacuum to provide 2 g of a black syrup. This material was purified
20 using silica gel column chromatography to provide 1.4 g of 2-butylthiazolo[4,5-c][1,5]naphthyridine as an amber liquid. High resolution mass spec (EI): Calculated for C₁₃H₁₃N₃S (M⁺) 243.0830; Found 243.0825

Part D


A solution of 3-chloroperoxybenzoic acid (1.1 g of 57-86%) in chloroform (50 mL)
25 was added in a steady stream to a solution of 2-butylthiazolo[4,5-c][1,5]naphthyridine (1.4 g, 5.8 mmol) in chloroform (100 mL). The reaction mixture was stirred at ambient temperature for 2.5 hours and then it was diluted with dichloromethane, washed twice with 10% sodium hydroxide, washed with brine, dried and concentrated under vacuum to provide a light yellow syrup which solidified on standing. This material was purified by
30 silica gel column chromatography to provide 1.2 g of a pale yellow solid. This material was recrystallized from petroleum ether (15 mL) and hexanes (100 mL) to provide 2-

butylthiazolo[4,5-*c*][1,5]naphthyridine-5N-oxide, m.p. 65-69°C. Analysis: Calculated for $C_{13}H_{13}N_3OS$: %C, 60.21; %H, 5.05; %N, 16.20; Found: %C, 60.43; %H, 5.17; %N, 16.18. High resolution mass spec (EI): Calculated for $C_{13}H_{13}N_3OS(M^+)$ 259.0779; Found 259.0789.

5

Example 58


2-Butylthiazolo[4,5-*c*][1,5]naphthyridin-4-amine



A solution of 2-butylthiazolo[4,5-*c*][1,5]naphthyridine-5N-oxide (0.5 g, 1.9 mmol) in dichloromethane (100 mL) was cooled in an ice bath. A solution of trichloroacetyl isocyanate (0.4 g, 2.1 mmol) in dichloromethane (25 mL) was added dropwise. The reaction mixture was stirred at ambient temperature for 8 hours. Added an amount of ammonia in methanol sufficient to make the reaction mixture basic and then let stand overnight. The reaction mixture was diluted with additional dichloromethane and then washed twice with 10% sodium hydroxide, washed with brine, dried and concentrated under vacuum to provide 0.6 g of a pale yellow solid. This material was purified by silica gel column chromatography and then recrystallized from acetonitrile (8 mL) to provide 0.15 g of 2-butylthiazolo[4,5-*c*][1,5]naphthyridin-4-amine as a white crystalline solid, m.p. 136-138°C. Analysis: Calculated for $C_{13}H_{14}N_4S$: %C, 60.44; %H, 5.46; %N, 21.69; Found: %C, 60.12; %H, 5.42; %N, 21.51. High resolution mass spec (EI) Calculated for $C_{13}H_{14}N_4S(M^+)$ 258.0941 Found: 258.0939. NMR chemical shifts in $CDCl_3$ (ppm) 8.637 dd (1H, $J=3.6; 1.2$ Hz), 8.048 dd (1H, $J=8.5; 1.2$ Hz), 7.486 dd (1H, $J=8.5; 3.6$ Hz), 5.691 bs(2H), 3.196 t (2H, $J=7$ Hz), 1.918 quintet (2H, $J=7$ Hz), 1.509 sextet (2H, $J=7$ Hz), 1.003 t (3H, $J=7$ Hz).

25

Example 59

2-Propylthiazolo[4,5-*c*][1,5]naphthyridine-5N-oxide5 Part A

Using the general method of Example 57 Part B, 3-amino[1,5]naphthyridin-4-ol (1.8 g, 11.2 mmol) was reacted with butyryl chloride (1.3 g, 12.3 mmol) to provide 1.2 g of N-(4-hydroxy[1,5]naphthyridin-3-yl)butanamide as a charcoal gray solid, m.p. >360°C.

Part B


10 Using the general method of Example 57 Part C, N-(4-hydroxy[1,5]naphthyridin-3-yl)butanamide (1.2 g, 5.2 mmol) was reacted with phosphorous pentasulfide (2.3 g, 5.2 mmol) to provide 0.9 g of 2-propylthiazolo[4,5-*c*][1,5]naphthyridine as an amber syrup.

Part C

15 Using the general method of Example 57 Part D, 2-propylthiazolo[4,5-*c*][1,5]naphthyridine (0.9 g, 3.9 mmol) was oxidized to provide 0.7 g of 2-propylthiazolo[4,5-*c*][1,5]naphthyridin-5N-oxide as a pale yellow solid, m.p. 139-142°C. Analysis: Calculated for C₁₂H₁₁N₃OS: %C, 58.76; %H, 4.52; %N, 17.13; Found: %C, 58.66; %H, 4.59; %N, 17.16. High resolution mass spec: (EI) calculated for C₁₂H₁₁N₃OS (M⁺) 245.0623; Found 245.0612.

20


Example 60

2-Propylthiazolo[4,5-*c*][1,5]naphthyridin-4-amine

Using the general method of Example 58, 2-propylthiazolo[4,5-c][1,5]naphthyridin-5N-oxide (0.5 g, 2 mmol) was aminated to provide 0.2 g of 2-propylthiazolo[4,5-c][1,5]naphthyridin-4-amine as ivory needles, m.p. 135-136°C. Analysis: Calculated for C₁₂H₁₂N₄S: %C, 58.99; %H, 4.95; %N, 22.93; Found: %C, 59.06; %H, 4.96; %N, 22.97. High resolution mass spec (EI) Calculated for C₁₂H₁₂N₄S (M⁺) 244.0783; Found 244.0785.

Example 61

2-Propylpyrido[3,4-d][1,3]thiazole-5N-oxide



Part A

A suspension of 3-nitropyridin-4-ol (1.0 g, 7.1 mmol) in methanol (110 mL) and a small amount of raney nickel catalyst were combined in a Parr bottle and hydrogenated for 4 hours. The reaction mixture was acidified with a solution of hydrochloric acid in ethanol and then filtered to remove catalyst. The filtrate was refiltered using Celite filter aid. The filtrate was concentrated under vacuum to provide 1.2 g of 3-aminopyridin-4-ol as a brown powder, m.p. 199-200°C.

Part B

N,N-diisopropylethylamine (33 mL, 180 mmol) was added to a suspension of 3-aminopyridin-4-ol (8.5 g, 46 mmol) in dichloromethane (100 mL). A solution of butyryl chloride (5.4 g, 51 mmol) in dichloromethane (100 mL) was added dropwise. The reaction mixture was stirred at ambient temperature for 3 hours and then at reflux for 3 hours. The reaction mixture was filtered to remove a black precipitate. The filtrate was concentrated under vacuum. The resulting light brown residue was triturated with hot ethyl acetate (250 mL) and then allowed to cool overnight. The mixture was filtered to remove solids (9.1 g) and the solids were washed with fresh ethyl acetate. The filtrate was concentrated under vacuum to provide 13 g of a light amber syrup. The syrup was dissolved in water and then extracted twice with ethyl acetate. The extracts were combined, washed with brine, dried

and then concentrated under vacuum to provide 2.5 g of an amber syrup. This material was purified by column chromatography to provide 1.2 g of N-(4-hydroxypyrid-3-yl)butanamide as a light amber syrup which solidified on standing.

Part C

- 5 Using the general method of Example 57 Part C, N-(4-hydroxypyrid-3-yl)butanamide (1.1 g, 6.1 mmol) was reacted with phosphorous pentasulfide (2.7 g, 6.1 mmol) to provide 0.4 g of 2-propylpyrido[3,4-d][1,3]thiazole as an amber syrup, which solidified on standing, mp 44-47°C.


Part D

- 10 Using the general method of Example 57 Part D, 2-propylpyrido[3,4-d][1,3]thiazole (0.4 g, 2.2 mmol) was oxidized to provide 0.2 g of 2-propylpyrido[3,4-d][1,3]thiazol-5N-oxide as short ivory needles after recrystallization from ethyl acetate (7 mL), m.p.137-139°C. Analysis: Calculated for C₉H₁₀N₂OS: %C, 55.65; %H, 5.19; %N, 14.42; Found: %C, 55.47; %H, 5.25; %N, 14.34.

15

Example 62

2-Propylpyrido[3,4-d][1,3]thiazol-4-amine Trifluoroacetate




- 20 A solution of trichloroacetyl isocyanate (0.11 g, 0.6 mmol) in dichloromethane (5 mL) was added dropwise to a chilled (ice bath) solution of 2-propylpyrido[3,4-d][1,3]thiazol-5N-oxide (0.1 g, 0.5 mmol) in dichloromethane (20 mL). The reaction mixture was stirred at ambient temperature for 5 hours. Additional trichloroacetyl isocyanate (0.2 g) was added and the reaction mixture was stirred at ambient temperature
- 25 overnight. The reaction mixture was briefly warmed to reflux then allowed to stir at ambient temperature for about 3 hours. Ammonia was bubbled into the reaction mixture which was then allowed to stir at ambient temperature for 1 hour. The reaction mixture was diluted with dichloromethane, washed twice with 10% sodium hydroxide, washed

with brine, dried and then concentrated under vacuum to provide an amber syrup. The reaction was repeated on the same scale. The products were combined to provide 0.1 g of an amber syrup. This material was purified by semi-preparative HPLC on a Gilson system (Rainin Microsorb C18 column, 21.4 x 250 mm, 8 micron particle size, 60A pore, 10 mL/min., gradient elution from 2-95% B in 25 min., hold at 95% B for 5 min., where A=0.1 % trifluoroacetic acid/water and B=0.1% trifluoroacetic acid/acetonitrile, peak detection at 254 nm for triggering fraction collection). The semi-prep HPLC fractions were analyzed by LC-APCI/MS and the appropriate fractions were lyophilized to provide the desired product as a trifluoroacetate salt m.p. 160-162°C. Analysis: Calculated for $C_9H_{11}N_3S + CF_3C(O)_2H$: %C, 42.99; %H, 3.94; %N, 13.67; Found: %C, 42.84; %H, 3.98; %N, 13.52. High resolution mass spec: (EI) calculated for $C_9H_{11}N_3S$ (M^+) 193.0674; Found 193.0681.

Example 63

7-Chloro-2-propylthiazolo[4,5-c]quinolin-4-amine



Part A

7-Chloro-4-hydroxyquinoline (35 g, 0.195 mol; available from Aldrich, Milwaukee, WI) and nitric acid (350 mL of 70%) were combined and heated at reflux for 75 minutes. The reaction mixture was poured over ice while still hot. The resulting bright yellow precipitate was isolated by filtration and then washed 3 times with boiling ethyl acetate to provide 17.3 g of 7-chloro-3-nitro-4-hydroxyquinoline as a pale yellow solid.

Part B

7-Chloro-3-nitro-4-hydroxyquinoline (4.48 g, 20 mmol), tin (II) chloride dihydrate (22.6 g, 100 mmol) and ethanol (200 mL) were combined and then heated at reflux for 4 hours. The reaction mixture was cooled to ambient temperature and then poured into

water (250 mL). The mixture was brought to neutral pH by the addition of saturated sodium bicarbonate and then filtered to remove tin salts. The filtrate was extracted with ethyl acetate. The combined organic fractions were dried over magnesium sulfate, filtered and then concentrated under vacuum to provide 1.8 g of 3-amino-7-chloro-4-hydroxyquinoline as a green powder.

Part C

Under a nitrogen atmosphere, butyryl chloride (0.76 mL, 7.3 mmol) was added dropwise to a mixture of 3-amino-7-chloro-4-hydroxyquinoline (1.3 g, 6.7 mmol), triethylamine (3.0 mL, 21.5 mmol) and anhydrous tetrahydrofuran (20 mL). The reaction mixture was maintained at ambient temperature overnight. The resulting precipitate was isolated by filtration, washed with water followed by tetrahydrofuran, and then vacuum dried to provide 1.05 g of N-(7-chloro-4-hydroxyquinolin-3-yl)butanamide as a tan powder.

Part D

A mixture of N-(7-chloro-4-hydroxyquinolin-3-yl)butanamide (0.9 g, 3.4 mol), phosphorous pentasulfide (1.51 g, 3.4 mmol) and pyridine (25 mL) was refluxed under a nitrogen atmosphere for 2.5 hours and then allowed to cool to ambient temperature. The reaction mixture was partitioned between dichloromethane (100 mL) and saturated sodium bicarbonate (100 mL). The aqueous layer was extracted with dichloromethane (2 X 100 mL). The organic fractions were combined, washed with water, dried over magnesium sulfate, filtered and then concentrated under vacuum to provide crude product. This material was purified by silica gel chromatography, (97:3 dichloromethane:methanol, 10 g SiO₂) to provide 0.62 g of 7-chloro-2-propylthiazolo[4,5-c]quinoline as a golden yellow solid.

Part E

Under a nitrogen atmosphere 3-chloroperoxybenzoic acid (0.7 g of 57-86%) was added to a mixture of 7-chloro-2-propylthiazolo[4,5-c]quinoline (0.5 g, 1.9 mmol) and chloroform (20 mL). After 2 hours at ambient temperature additional 3-chloroperoxybenzoic acid (0.2 g) was added and the reaction mixture was maintained at ambient temperature for 14 hours. The reaction mixture was diluted with dichloromethane and then washed twice with saturated sodium bicarbonate. The organic fraction was dried


over magnesium sulfate, filtered and then concentrated under vacuum to provide 0.52 g of 7-chloro-2-propylthiazolo[4,5-*c*]quinoline-5N-oxide as an orange solid.

Part F

Under a nitrogen atmosphere trichloroacetyl isocyanate (0.32 mL, 2.7 mmol) was added to a mixture of 7-chloro-2-propylthiazolo[4,5-*c*]quinoline-5N-oxide (0.50 g, 1.8 mmol) and dichloromethane (20 mL). The reaction mixture was maintained at ambient temperature for 2 hours and then concentrated under vacuum. The resulting oily residue was dissolved in methanol (10 mL), sodium methoxide (1 mL of 25%, 4.4 mmol) was added and the reaction mixture was maintained at ambient temperature for 2.5 days. The resulting precipitate was isolated by filtration and washed with hexane to provide 0.28 g of the desired product as a golden yellow powder. A 50 mg portion was recrystallized from methanol to provide 7-chloro-2-propylthiazolo[4,5-*c*]quinolin-4-amine as a golden yellow crystalline solid, m.p. 159-160°C. ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.82 (d, *J*=8.5 Hz, 1H), 7.60 (d, *J*=2.0 Hz, 1H), 7.27 (dd, *J*=8.5, 2.1 Hz, 1H), 7.10 (s, 2H), 3.16 (t, *J*=7.4 Hz, 2H), 1.87 (sextet, *J*=7.4 Hz, 2H), 1.02 (t, *J*=7.4 Hz, 3H); MS (EI) *m/e* 277.0441 (277.0440 calcd for C₁₃H₁₂ClN₃S).

Example 64

7-Methoxy-2-propylthiazolo[4,5-*c*]quinolin-4-amine



Part A

3-Methoxyaniline (12.3 g, 0.1 mol) and diethyl ethoxymethylenemalonate (21.6 g, 0.1 mol) were combined and heated at 120°C for 3 hours. The reaction mixture was allowed to cool to ambient temperature and then placed under vacuum overnight to provide 28.5 g of diethyl 2-[3-(methoxyanilino)methylene]malonate as an orange oil.

Part B

Dowtherm A (~ 200 mL) was charged into a flask equipped with a stir bar, nitrogen inlet, Dean-Stark trap and condenser. The solvent was heated to a vigorous reflux and then 2-[3-(methoxyanilino)methylene]malonate (20.0 g, 68 mmol) was added. The reaction mixture was heated for 0.5 hr. and the brown solution was allowed to cool to ambient temperature. The resulting precipitate was isolated by filtration, washed with acetone and then air dried to provide 12.5 g of ethyl 4-hydroxy-7-methoxyquinoline-3-carboxylate as a yellow powder.

Part C

A suspension of ethyl 4-hydroxy-7-methoxyquinoline-3-carboxylate (12.0 g, 48 mmol) in 10% sodium hydroxide/water (200 mL) was heated at reflux for 1.5 hr. The reaction mixture was allowed to cool to ambient temperature and then made acidic (pH = 3) by the dropwise addition of concentrated hydrochloric acid. The resulting precipitate was isolated by filtration, washed twice with water and then dried overnight in a vacuum oven at 80°C to provide 10.4 g of 4-hydroxy-7-methoxyquinoline-3-carboxylic acid.

Part D

A suspension of 4-hydroxy-7-methoxyquinoline-3-carboxylic acid (4.0 g) in Dowtherm A (75 mL) was heated at reflux for 2 hrs. The resulting brown solution was allowed to slowly cool to ambient temperature. The resulting precipitate was isolated by filtration and then dried in a vacuum oven at 80°C for 2.5 days to provide 3.1 g of 7-methoxyquinolin-4-ol as a light tan solid.

Part E

A mixture of 7-methoxyquinolin-4-ol (5.0 g, 28.5 mmol) and propionic acid (50 mL) was heated to reflux. Nitric acid (3.2 mL of 70%, 50 mmol) was added dropwise over a period of 15 minutes. The reaction mixture was refluxed for 2 hrs and then allowed to cool to ambient temperature. The resulting precipitate was isolated by filtration, washed with cold ethanol followed by hexanes and then dried to provide 3.9 g of 7-methoxy-3-nitroquinolin-4-ol as a gray solid.

Part F

7-Methoxy-3-nitroquinolin-4-ol (4.5 g, 20.4 mmol), methanol (250 mL), ammonium hydroxide (5 mL) and palladium on carbon (400 mg of 10%) were combined. The mixture was placed on a Parr apparatus under a hydrogen atmosphere at 40 psi (2.8 Kg/cm²) for 2 hrs. The reaction mixture was filtered. The filtrate was concentrated under

vacuum to provide a green solid. This material was dissolved in methanol (20 mL) and then 1N hydrochloric acid in diethyl ether (75 mL) was added. The resulting precipitate was isolated by filtration and dried to provide 2.6 g of 3-amino-7-methoxyquinolin-4-ol hydrochloride as a pink solid.

5 Part G

Butyryl chloride (0.63 mL, 6.1 mmol) was added dropwise to a solution containing 3-amino-7-methoxyquinolin-4-ol hydrochloride (1.0 g, 5.26 mmol), triethylamine (2.35 mL, 16.8 mmol), dichloromethane (30 mL) and N,N-dimethylformamide (10 mL). The reaction mixture was maintained at ambient temperature overnight. The N,N-dimethylformamide was removed under vacuum and the resulting solid was partitioned between dichloromethane (100 mL) and water (100 mL). The organic fraction was washed with water, dried over magnesium sulfate, filtered and then concentrated under vacuum to provide 0.86 g of N-(4-hydroxy-7-methoxyquinolin-3-yl)butanamide as a tan solid.

15 Part H

Under a nitrogen atmosphere a mixture of N-(4-hydroxy-7-methoxyquinolin-3-yl)butanamide (0.66 g, 2.54 mmol), pyridine (20 mL), and phosphorous pentasulfide (1.13 g, 2.54 mmol) was heated at reflux and then cooled to ambient temperature. The reaction mixture was filtered. The filtrate was partitioned between dichloromethane (100 mL) and saturated sodium bicarbonate (100 mL). The aqueous fraction was extracted with additional dichloromethane (100 mL). The organic fractions were combined, washed with water, dried over magnesium sulfate, filtered and then concentrated under vacuum to provide a solid. This material was purified by silica gel chromatography (15 g of SiO₂ eluting with 95:5 dichloromethane:methanol) to provide 0.45 g of 7-methoxy-2-propylthiazolo[4,5-c]quinoline as a pale yellow powder.

25 Part I

Using the method of Example 63 Part E, 7-methoxy-2-propylthiazolo[4,5-c]quinoline (0.40 g, 1.55 mmol) was oxidized to provide 7-methoxy-2-propylthiazolo[4,5-c]quinoline-5N-oxide as an orange solid.

30 Part J

Using the method of Example 63 Part F, the N-oxide from Part I was reacted with trichloroacetyl isocyanate and the resulting amide was hydrolyzed to provide 190 mg of the desired product as an off white solid. An analytical sample was obtained by recrystallization from methanol to provide 7-methoxy-2-propylthiazolo[4,5-c]quinolin-4-

amine as off-white needles, m.p. 152-154°C. ¹H NMR (300 MHz, DMSO-d₆) δ 7.67 (d, J=8.7 Hz, 1H), 7.06 (d, J=2.4 Hz, 1H), 6.91 (dd, J=8.7, 2.5 Hz, 1H), 6.82 (s, 2H), 3.86 (s, 3H), 3.11 (t, J=7.4 Hz, 2H), 1.85 (sextet, J=7.4 Hz, 2H), 1.01 (t, J=7.4 Hz, 3H); MS (EI) m/e 273.0934 (273.0936 calcd for C₁₄H₁₅N₃OS).

5

INTERFERON (α) INDUCTION IN HUMAN CELLS

An in vitro human blood cell system was used to assess interferon induction by compounds of the invention. Activity is based on the measurement of interferon secreted into culture media. Interferon is measured by bioassay.

10

Blood Cell Preparation for Culture

Whole blood is collected by venipuncture into EDTA vacutainer tubes. Peripheral blood mononuclear cells (PBM's) are separated from whole blood by using either LeucoPREP™ Brand Cell Separation Tubes (available from Becton Dickinson) or Ficoll-Paque® solution (available from Pharmacia LKB Biotechnology Inc, Piscataway, NJ). The PBM's are suspended at 1 x 10⁶/mL in RPMI 1640 media (available from GIBC(O), Grand Island, NY) containing 25 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid) and L-glutamine (1% penicillin-streptomycin solution added) with 10% heat inactivated (56°C for 30 minutes) fetal calf serum added. 200 μL portions of PBM suspension are added to 96 well (flat bottom) sterile tissue culture plates (Becton Dickinson Labware, Lincoln Park, NJ).

15

20

Compound Preparation

The compounds are solubilized in ethanol, dimethyl sulfoxide or tissue culture water then diluted with tissue culture water, 0.01N sodium hydroxide or 0.01N hydrochloric acid (The choice of solvent will depend on the chemical characteristics of the compound being tested.). Ethanol or DMSO concentration should not exceed a final concentration of 1% for addition to the culture wells. Compounds are initially tested in a concentration range of from about 0.1 μg/mL to about 5 μg/mL. Compounds which show induction at a concentration of 0.5 μg/mL are then tested in a wider concentration range.

25

30

Incubation

The solution of test compound is added in a volume (less than or equal to 50 μ L) to the wells containing 200 μ L of diluted whole blood or of PBM's in media. Solvent and/or media is added to control wells (wells with no test compound) and as needed to adjust the final volume of each well to 250 μ L. The plates are covered with plastic lids, vortexed gently and then incubated for 48 hours at 37°C with a 5% carbon dioxide atmosphere.

Separation

Following incubation, the plates are covered with parafilm and then centrifuged at 1000 rpm for 10 to 15 minutes at 4°C in a Damon IEC Model CRU-5000 centrifuge. Media (about 200 μ L) is removed from 4 to 8 wells and pooled into 2 mL sterile freezing vials. Samples are maintained at -70°C until analysis.

Interferon Analysis/Calculation

Interferon is determined by bioassay using A549 human lung carcinoma cells challenged with encephalomyocarditis. The details of the bioassay method have been described by G. L. Brennan and L. H. Kronenberg in "Automated Bioassay of Interferons in Micro-test Plates", Biotechniques, June/July, 78, 1983, incorporated herein by reference. Briefly stated the method is as follows: interferon dilutions and A549 cells are incubated at 37°C for 12 to 24 hours. The incubated cells are infected with an inoculum of encephalomyocarditis virus. The infected cells are incubated for an additional period at 37°C before quantifying for viral cytopathic effect. The viral cytopathic effect is quantified by staining followed by spectrophotometric absorbance measurements. Results are expressed as alpha reference units/mL based on the value obtained for NIH HU IF-L standard. The interferon was identified as essentially all interferon alpha by testing in checkerboard neutralization assays against rabbit anti-human interferon (beta) and goat anti-human interferon (alpha) using A549 cell monolayers challenged with encephalomyocarditis virus.

Compounds of the invention were tested for their ability to induce interferon in human cells using the test method described above. The results are given in the table below where "+" indicates that the compound induced interferon α at that particular concentration, a "-" indicates that the compound did not induce interferon α at that

particular concentration, and a “±” indicates that the results were equivocal at that particular concentration.

Interferon (α) Induction in Human Cells												
Example	Dose Concentration ($\mu\text{g/mL}$)											
	0.01	0.05	0.10	0.50	1.0	5.0	10.0	25.0	50.0			
2	-	-	±	+	+	+	not run	not run	not run			
4	-	-	-	+	+	+	not run	not run	not run			
5	-	-	-	-	+	+	+	+	+			
8	-	-	-	+	+	+	not run	not run	not run			
12	-	+	+	+	+	+	not run	not run	not run			
14	-	-	-	+	+	+	not run	not run	not run			
16	-	-	-	-	+	+	not run	not run	not run			
18	-	-	+	+	+	+	not run	not run	not run			
20	-	-	-	-	-	+	+	+	+			
22	-	-	-	-	-	-	+	+	+			
24	-	-	-	-	-	-	+	+	+			
26	-	-	-	-	-	+	+	+	+			
28	-	-	-	+	+	+	not run	not run	not run			
30	-	-	-	-	+	+	not run	not run	not run			
32	-	-	±	+	+	+	not run	not run	not run			
34	-	-	-	-	-	-	not run	not run	not run			

Interferon (α) Induction in Human Cells												
Example	Dose Concentration ($\mu\text{g/mL}$)											
	0.01	0.05	0.10	0.50	1.0	5.0	10.0	25.0	50.0			
36	-	-	-	-	+	+	not run	not run	not run			
37	-	-	-	-	-	-	not run	not run	not run			
38	-	-	-	+	+	+	not run	not run	not run			
39	-	-	+	+	+	+	not run	not run	not run			
41	+	+	+	+	+	+	not run	not run	not run			
42	-	-	-	+	+	+	not run	not run	not run			
43	-	+	+	+	+	+	not run	not run	not run			
44	not run	not run	-	-	-	+	+	+	+			
45	-	+	+	+	+	+	not run	not run	not run			
46	-	+	+	+	+	+	not run	not run	not run			
47	-	+	+	+	+	+	not run	not run	not run			
48	-	-	-	+	+	+	not run	not run	not run			
49	-	-	-	+	+	+	not run	not run	not run			
50	not run	not run	-	+	+	+	+	+	+			
53	-	-	-	-	-	-	-	+	+			
54	-	-	-	-	-	-	-	+	+			

CYTOKINE INDUCTION IN HUMAN CELLS

An in vitro human blood cell system was used to assess cytokine induction by compounds of the invention. Activity is based on the measurement of interferon (α) and tumor necrosis factor (α) (IFN and TNF, respectively) secreted into culture media as described by Testerman et. al. In "Cytokine Induction by the Immunomodulators Imiquimod and S-27609", Journal of Leukocyte Biology, **58**, 365-372 (September, 1995).

Blood Cell Preparation for Culture

Whole blood is collected by venipuncture into EDTA vacutainer tubes from healthy human donors. Peripheral blood mononuclear cells (PBMCs) are separated from whole blood by density gradient centrifugation using Histopaque®-1077 (Sigma Chemicals, St. Louis, MO). The PBMCs are suspended at $3-4 \times 10^6$ cells/mL in RPMI 1640 medium containing 10 % fetal bovine serum, 2 mM L-glutamine and 1% penicillin/streptomycin solution (RPMI complete). The PBMC suspension is added to 48 well flat bottom sterile tissue culture plates (Costar, Cambridge, MA or Becton Dickinson Labware, Lincoln Park, NJ) containing an equal volume of RPMI complete media containing test compound.

Compound Preparation

The compounds are solubilized in dimethyl sulfoxide (DMSO). The DMSO concentration should not exceed a final concentration of 1% for addition to the culture wells. The compounds are generally tested in a concentration range of from 0.12 to 30 μ M.

Incubation

The solution of test compound is added at 60 μ M to the first well containing RPMI complete and serial 3 fold dilutions are made. The PBMC suspension is then added to the wells in an equal volume, bringing the test compound concentrations to the desired range (0.12 to 30 μ M). The final concentration of PBMC suspension is $1.5-2 \times 10^6$ cells/mL. The plates are covered with sterile plastic lids, mixed gently and then incubated for 18 to 24 hours at 37°C in a 5% carbon dioxide atmosphere.

Separation

Following incubation the plates are centrifuged for 5-10 minutes at 1000 rpm (~200 x g) at 4°C. The cell culture supernatant is removed with a sterile polypropylene pipet and transferred to sterile polypropylene tubes. Samples are maintained at -30 to -70°C until analysis. The samples are analyzed for interferon (α) by either ELISA or bioassay and for tumor necrosis factor (α) by ELISA

Interferon Bioassay Analysis

Interferon is determined by bioassay using A549 human lung carcinoma cells challenged with encephalomyocarditis. The details of the bioassay method have been described by G. L. Brennan and L. H. Kronenberg in "Automated Bioassay of Interferons in Micro-test Plates", Biotechniques, June/July, 78, (1983), incorporated herein by reference. Briefly stated the method is as follows: A549 cells are incubated with dilutions of samples or a standard interferon at 37°C for 24 hours. The incubated cells are then infected with an inoculum of encephalomyocarditis virus. The infected cells are incubated for an additional 24 hours at 37°C before evaluating for viral cytopathic effect. The viral cytopathic effect is quantified by staining with crystal violet followed by visual scoring of the plates. Results are expressed as alpha reference units/mL based on the value obtained for NIH Human Leukocyte IFN standard.

Interferon (α) and Tumor Necrosis Factor (α) Analysis by ELISA

Interferon (α) concentration is determined by ELISA using a Human Multi-Species kit from PBL Biomedical Laboratories, New Brunswick, NJ according to the manufacturer's instructions.

Tumor necrosis factor (α) (TNF) concentration is determined using ELISA kits available from Genzyme, Cambridge, MA; R&D Systems, Minneapolis, MN; or Pharmingen, San Diego, CA according to the manufacturer's instructions.

In the tables below, a "+" indicates that the compound induced the indicated cytokine at that particular concentration, a "-" indicates that the compound did not induce the indicated cytokine at that particular concentration, and a "±" indicates that the results were equivocal at that particular concentration.

5

Cytokine Induction in Human Cells														
Example	Interferon (α) Induction						Tumor Necrosis Factor Induction							
	Dose Concentration (μ M)						Dose Concentration (μ M)							
	0.12	0.37	1.11	3.33	10	30	0.12	0.37	1.11	3.33	10	30		
12	-	-	-	+	+	+	+	+	+	+	+	+		
18	-	-	+	+	+	+	+	+	+	+	+	+		
20	-	-	+	-	-	-	-	-	-	+	+	+		
24	-	-	-	-	-	-	-	-	-	-	-	+		
42	-	-	-	-	+	-	-	-	+	+	+	+		
53	-	-	-	-	-	-	-	-	-	-	-	-		
58	-	-	-	-	\pm	\pm	-	+	+	+	+	+		
60	-	-	-	-	-	-	-	\pm	+	+	+	+		
62	-	-	-	-	-	-	-	-	-	-	-	-		
63	-	-	+	+	+	-	+	+	+	+	+	+		
64	-	+	+	+	+	+	+	+	+	+	+	+		

The present invention has been described with reference to several embodiments thereof. The foregoing detailed description and examples have been provided for clarity of understanding only, and no unnecessary limitations are to be understood therefrom. It will be apparent to those skilled in the art that many changes can be made to the described
5 embodiments without departing from the spirit and scope of the invention. Thus, the scope of the invention should not be limited to the exact details of the compositions and structures described herein, but rather by the language of the claims that follow.


10

9

WE CLAIM:

1. A compound of the formula I:

5



(I)

wherein:

R_1 is selected from the group consisting of oxygen, sulfur and selenium;

R_2 is selected from the group consisting of

10

-hydrogen;

-alkyl;

-alkyl-OH;

-haloalkyl;

-alkenyl;

15

-alkyl-X-alkyl;

-alkyl-X-alkenyl;

-alkenyl-X-alkyl;

-alkenyl-X-alkenyl;

-alkyl-N(R_5)₂;

20

-alkyl-N₃;

-alkyl-O-C(O)-N(R_5)₂;

-heterocyclyl;

-alkyl-X-heterocyclyl;

-alkenyl-X-heterocyclyl;

25

-aryl;

-alkyl-X-aryl;
-alkenyl-X-aryl;
-heteroaryl;
-alkyl-X-heteroaryl; and
5 -alkenyl-X-heteroaryl;

R_3 and R_4 are each independently:

-hydrogen;
-X-alkyl;
-halo;
10 -haloalkyl;
-N(R_5)₂;

or when taken together, R_3 and R_4 form a fused
aromatic, heteroaromatic, cycloalkyl or heterocyclic ring;

X is selected from the group consisting of $-O-$, $-S-$, $-NR_5-$, $-C(O)-$, $-C(O)O-$,
15 $-OC(O)-$, and a bond; and

each R_5 is independently H or C₁₋₈alkyl;

with the proviso that when R_1 is sulfur, R_3 is not $-NH_2$; or a pharmaceutically
acceptable salt thereof.

20 2. A compound according to claim 1 wherein R_1 is oxygen or sulfur.

3. A compound according to claim 1, wherein R_3 and R_4 are taken together to
form a substituted or unsubstituted benzene ring.

25 4. A compound according to claim 2 wherein R_3 and R_4 are taken together to
form a substituted or unsubstituted benzene ring.

5. A compound according to claim 1 wherein R_3 and R_4 are taken together to
form a substituted or unsubstituted pyridine ring.

30

6. A compound according to claim 2 wherein R₃ and R₄ are taken together to form a substituted or unsubstituted pyridine ring.

7. A compound according to claim 1 wherein R₂ is C₁₋₄ alkyl.

5

8. A compound according to claim 2 wherein R₂ is C₁₋₄ alkyl.

9. A compound according to claim 1 wherein R₁ is sulfur.

10

10. A compound according to claim 9 wherein R₃ and R₄ are taken together to form a substituted or unsubstituted benzene ring.

11. A compound according to claim 10 wherein R₂ is C₁₋₄ alkyl.

15

12. A compound according to claim 10 wherein R₂ is *n*-propyl.

13. 2-*n*-propylthiazolo[4,5-*c*]quinolin-4-amine, or a pharmaceutically acceptable salt thereof.

20

14. A compound selected from the group consisting of:

2-methylthiazolo[4,5-*c*]quinolin-4-amine;

thiazolo[4,5-*c*]quinolin-4-amine;

2-ethylthiazolo[4,5-*c*]quinolin-4-amine;

2-propylthiazolo[4,5-*c*]quinolin-4-amine;

25

2-pentylthiazolo[4,5-*c*]quinolin-4-amine;

2-butylthiazolo[4,5-*c*]quinolin-4-amine;

2-(1-methylethyl)thiazolo[4,5-*c*]quinolin-4-amine;

2-(2-phenyl-1-ethenyl)thiazolo[4,5-*c*]quinolin-4-amine;

2-(2-phenyl-1-ethyl)thiazolo[4,5-*c*]quinolin-4-amine;

30


2-(4-aminothiazolo[4,5-*c*]quinolin-2-yl)-1,1-dimethylethyl carbamate;

2-(ethoxymethyl)thiazolo[4,5-*c*]quinolin-4-amine;

2-(methoxymethyl)thiazolo[4,5-c]quinolin-4-amine;
2-(2-methylpropyl)thiazolo[4,5-c]quinolin-4-amine;
2-benzylthiazolo[4,5-c]quinolin-4-amine;
8-methyl-2-propylthiazolo[4,5-c]quinolin-4-amine;
5 (4-aminothiazolo[4,5-c]quinolin-2-yl)methanol;
2-methyloxazolo[4,5-c]quinolin-4-amine;
2-ethyloxazolo[4,5-c]quinolin-4-amine;
2-butyloxazolo[4,5-c]quinolin-4-amine;
2-propylthiazolo [4,5-c] quinolin-4,8-diamine;
10 2-propyloxazolo [4,5-c] quinolin-4-amine;
8-bromo-2-propylthiazolo[4,5-c]quinolin-4-amine;
7-methyl-2-propylthiazolo[4,5-c]quinolin-4-amine;
2-butyl-7-methyloxazolo[4,5-c]quinolin-4-amine;
7-methyl-2-propyloxazolo[4,5-c]quinolin-4-amine;
15 7-fluoro-2-propyloxazolo[4,5-c]quinolin-4-amine;
7-fluoro-2-propylthiazolo[4,5-c]quinolin-4-amine;
2-propyl-7-(trifluoromethyl)thiazolo[4,5-c]quinolin-4-amine;
2-(4-morpholino)thiazolo[4,5-c]quinolin-4-amine;
2-(1-pyrrolidino)thiazolo[4,5-c]quinolin-4-amine;
20 2-butylthiazolo[4,5-c][1,5]naphthyridin-4-amine;
2-propylthiazolo[4,5-c][1,5]naphthyridin-4-amine;
7-chloro-2-propylthiazolo[4,5-c]quinolin-4-amine;
7-methoxy-2-propylthiazolo[4,5-c]quinolin-4-amine;
and pharmaceutically acceptable salts thereof.

25

15. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula I(a):



I(a)

wherein:

R_1 is selected from the group consisting of oxygen, sulfur and selenium;

R_2 is selected from the group consisting of

- 5 -hydrogen;
- alkyl;
- alkyl-OH;
- haloalkyl;
- alkenyl;
- 10 -alkyl-X-alkyl;
- alkyl-X-alkenyl;
- alkenyl-X-alkyl;
- alkenyl-X-alkenyl;
- alkyl-N(R_5)₂;
- 15 -alkyl-N₃;
- alkyl-O-C(O)-N(R_5)₂;
- heterocyclyl;
- alkyl-X-heterocyclyl;
- alkenyl-X-heterocyclyl;
- 20 -aryl;
- alkyl-X-aryl;
- alkenyl-X-aryl;
- heteroaryl;
- alkyl-X-heteroaryl; and
- 25 -alkenyl-X-heteroaryl;

R_3 and R_4 are each independently:

-hydrogen;

-X-alkyl;

-halo;

-haloalkyl;

5 -N(R₅)₂;

or when taken together, R₃ and R₄ form a fused

aromatic, heteroaromatic, cycloalkyl or heterocyclic ring;

X is selected from the group consisting of -O-, -S-, -NR₅-, -C(O)-, -C(O)O-,
-OC(O)-, and a bond; and

10 each R₅ is independently H or C₁₋₈alkyl; or a pharmaceutically acceptable salt
thereof, with a pharmaceutically acceptable carrier.

16. A method of inducing cytokine biosynthesis in a mammal comprising
administering a composition of claim 15 to the mammal.

15

17. The method of claim 16 wherein the cytokine comprises IFN- α .

18. The method of claim 16 wherein the cytokine comprises TNF- α .

20 19. The method of claim 16 wherein the composition is administered topically.


20. A method of treating a viral disease in a mammal comprising administering
a composition of claim 15 to the mammal.

25 21. The method of claim 20 wherein the composition is administered topically.

22. A method of treating a neoplastic disease in a mammal comprising
administering a composition of claim 15 to the mammal.

30 23. The method of claim 22 wherein the composition is administered topically.

24. A compound of the formula II:



wherein

5 R_1 is selected from the group consisting of oxygen, sulfur and selenium;

R_2 is selected from the group consisting of

- hydrogen;
- alkyl;
- alkyl-OH;
- 10 -haloalkyl;
- alkenyl;
- alkyl-X-alkyl;
- alkyl-X-alkenyl;
- alkenyl-X-alkyl;
- 15 -alkenyl-X-alkenyl;
- alkyl-N(R₅)₂;
- alkyl-N₃;
- alkyl-O-C(O)-N(R₅)₂;
- heterocyclyl;
- 20 -alkyl-X-heterocyclyl;
- alkenyl-X-heterocyclyl;
- aryl;
- alkyl-X-aryl;
- alkenyl-X-aryl;
- 25 -heteroaryl;
- alkyl-X-heteroaryl;

-alkenyl-X-heteroaryl;

-SO₂CH₃; and

-CH₂-O-C(O)-CH₃;

R₃ and **R₄** are each independently:

5 -hydrogen;

-X-alkyl;

-halo;

-haloalkyl;

-N(R₅)₂;

10 or when taken together, **R₃** and **R₄** form a fused
aromatic, heteroaromatic, cycloalkyl or heterocyclic ring;

X is selected from the group consisting of -O-, -S-, -NR₅-, -C(O)-, -C(O)O-,
and a bond; and

each **R₅** is independently H or C₁₋₈alkyl.

15

25. A compound selected from the group consisting of:

2-methylthiazolo[4,5-*c*]quinoline-5N-oxide;

2-ethylthiazolo[4,5-*c*]quinoline-5N-oxide;

2-propylthiazolo[4,5-*c*]quinoline-5N-oxide;

20 2-pentylthiazolo[4,5-*c*]quinoline-5N-oxide;

2-butylthiazolo[4,5-*c*]quinoline-5N-oxide;

2-(1-methylethyl)thiazolo[4,5-*c*]quinoline-5N-oxide;

2-(2-phenyl-1-ethenyl)thiazolo[4,5-*c*]quinoline-5N-oxide;

2-phenylethylthiazolo[4,5-*c*]quinoline-5N-oxide;

25 2-methyl-1-thiazolo[4,5-*c*]quinolin-2-yl-2-propanol-5N-oxide;

2-(ethoxymethyl)thiazolo[4,5-*c*]quinoline-5N-oxide;

2-(methoxymethyl)thiazolo[4,5-*c*]quinoline-5N-oxide;

2-(2-methylpropyl)thiazolo[4,5-*c*]quinoline-5N-oxide;

2-benzylthiazolo[4,5-*c*]quinoline-5N-oxide;

30 8-methyl-2-propylthiazolo[4,5-*c*]quinoline-5N-oxide; and

2-butyloxazolo[4,5-*c*]quinoline-5N-oxide.

INTERNATIONAL SEARCH REPORT

Interr. Application No
PCT/US 99/17027

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D498/04 A61K31/44 C07D513/04 C07D517/04 C07D513/14
/(C07D498/04,263:00,221:00),(C07D513/04,277:00,221:00),
(C07D517/04,293:00,221:00),(C07D513/14,277:00,221:00,221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 16514 A (AMERICAN CYANAMID CO.) 23 April 1998 (1998-04-23) pages 1-3,9,12-19; claims ---	1,15-25
A	EP 0 055 248 A (AMERICAN HOME PRODUCTS CORP.) 30 June 1982 (1982-06-30) the whole document ---	1,15
A	US 4 038 396 A (TSUNG-YING SHEN) 26 July 1977 (1977-07-26) cited in the application the whole document ---	1,15
A	WO 93 05042 A (MINNESOTA MINING AND MANUFACTURING CO.) 18 March 1993 (1993-03-18) cited in the application the whole document ---	1,15-25
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

4 November 1999

Date of mailing of the international search report

19/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 99/17027

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 02598 A (MINNESOTA MINING AND MANUFACTURING CO.) 26 January 1995 (1995-01-26) cited in the application the whole document ---	1,15-25
A	WO 95 02597 A (MINNESOTA MINING AND MANUFACTURING CO.) 26 January 1995 (1995-01-26) cited in the application the whole document ---	1,15-25
A	TEMPLE C, JR. ET AL: "Synthesis of imidazo'4,5-b!pyridines and v-triazolo'4,5-b!pyridines. Preparation of 1-deaza-6-thioguanine analogs" J. ORG. CHEM. (JOCEAH);1976; VOL.41 (24); PP.3784-8, XP002121358 South. Res. Inst.;Kettering-Meyer Lab.; Birmingham; Ala. the whole document ---	1
P,A	WO 98 42712 A (SYNTHELABO) 1 October 1998 (1998-10-01)	1,15
P,X	pages 28,29, paragraph 4.1.4 -----	24

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 17027

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16-23
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 16-23
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: al Application No

PCT/US 99/17027

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9816514 A	23-04-1998	AU 4980697 A	11-05-1998
EP 55248 A	30-06-1982	US 4275065 A	23-06-1981
		AT 8787 T	15-08-1984
		AU 540269 B	08-11-1984
		AU 5943980 A	08-01-1981
		CA 1151170 A	02-08-1983
		CA 1151171 A	02-08-1983
		DK 266280 A	22-12-1980
		EP 0021773 A	07-01-1981
		ES 492671 A	01-06-1981
		GB 2055361 A,B	04-03-1981
		IE 50099 B	19-02-1986
		IE 50098 B	19-02-1986
		JP 56007793 A	27-01-1981
		KR 8400590 B	24-04-1981
		ZA 8003569 A	28-04-1982
		ZA 8101732 A	28-04-1982
		GB 2096995 A,B	27-10-1982
US 4038396 A	26-07-1977	US 4131677 A	26-12-1978
WO 9305042 A	18-03-1993	US 5268376 A	07-12-1993
		AU 2514792 A	05-04-1993
		CA 2116782 A	18-03-1993
		CZ 281726 B	11-12-1996
		CZ 9400487 A	13-07-1994
		EP 0603251 A	29-06-1994
		HU 67398 A	28-04-1995
		HU 69407 A	28-09-1995
		HU 9500663 A	28-11-1995
		IL 102951 A	30-09-1997
		JP 6510299 T	17-11-1994
		MX 9205046 A	01-03-1993
		NZ 244075 A	26-05-1995
		US 5525612 A	11-06-1996
		US 5714608 A	03-02-1998
		US 5346905 A	13-09-1994
		ZA 9206456 A	04-03-1993
WO 9502598 A	26-01-1995	US 5352784 A	04-10-1994
		EP 0708773 A	01-05-1996
		JP 9500628 T	21-01-1997
		US 5444065 A	22-08-1995
		US 5627281 A	06-05-1997
		US 5886006 A	23-03-1999
WO 9502597 A	26-01-1995	US 5366974 A	22-11-1994
		AU 693747 B	02-07-1998
		AU 4098797 A	08-01-1998
		AU 681687 B	04-09-1997
		AU 7247594 A	13-02-1995
		CZ 9600105 A	12-06-1996
		EP 0708772 A	01-05-1996
		IL 110091 A	18-03-1997
		IL 119546 A	04-01-1998
		IL 119547 A	04-01-1998
		JP 9500128 T	07-01-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/US 99/17027

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9502597 A		MX 9405383 A US 5446153 A US 5494916 A	31-01-1995 29-08-1995 27-02-1996
WO 9842712 A	01-10-1998	FR 2761071 A AU 6923998 A ZA 9802362 A	25-09-1998 20-10-1998 23-09-1998